

Musculoskeletal Rehabilitation

| ICD-10 CODE | DESCRIPTION | % OF CLINICAL GROUPING |
|-------------|--|------------------------|
| Z47.1 | Aftercare following joint replacement surgery | 21% |
| Z47.89 | Encounter for other orthopedic aftercare | 7% |
| M17.0 | Bilateral primary osteoarthritis of knee | 6% |
| M15.0 | Primary generalized (osteo)arthritis | 5% |
| Z47.81 | Encounter for orthopedic aftercare following surgical amp | 3% |
| S72.001D | Fx unsp part of nk of r femr, subs for clos fx w routn heal | 3% |
| I89.0 | Lymphedema, not elsewhere classified | 2% |
| S72.002D | Fx unsp part of nk of l femr, subs for clos fx w routn heal | 2% |
| S72.142D | Displ intertroch fx l femur, subs for clos fx w routn heal | 2% |
| M48.061 | Spinal stenosis, lumbar region without neurogenic claudication | 2% |
| S72.141D | Displ intertroch fx r femur, subs for clos fx w routn heal | 1% |
| M17.11 | Unilateral primary osteoarthritis, right knee | 1% |
| M54.16 | Radiculopathy, lumbar region | 1% |
| M51.36 | Other intervertebral disc degeneration, lumbar region | 1% |
| M17.12 | Unilateral primary osteoarthritis, left knee | 1% |
| M80.08XD | Age-rel osteopor w crnt path fx, verteb, 7thD | 1% |
| M54.2 | Cervicalgia | 1% |
| M15.9 | Polyosteoarthritis, unspecified | 1% |
| S32.010D | Wedge comprsn fx first lum vert, subs for fx w routn heal | 1% |
| M81.0 | Age-related osteoporosis w/o current pathological fracture | 1% |

TOP 20 DX BY CLINICAL GROUPING

Neuro/Stroke Rehabilitation

| ICD-10 CODE | DESCRIPTION | % OF CLINICAL GROUPING |
|-------------|--|------------------------|
| G20. | Parkinson's disease | 21% |
| I69.354 | Hemiplegia following cerebral infarction affecting left nondominant side | 13% |
| I69.351 | Hemiplegia following cerebral infarction affecting right dominant side | 13% |
| G30.9 | Alzheimer's disease, unspecified | 8% |
| G35. | Multiple sclerosis | 6% |
| I69.398 | Other sequelae of cerebral infarction | 5% |
| I69.320 | Aphasia following cerebral infarction | 2% |
| G12.21 | Amyotrophic lateral sclerosis | 2% |
| S06.5X0D | Traumatic subdural hemorrhage without loss of consciousness, substance | 1% |
| G31.83 | Dementia with Lewy bodies | 1% |
| G30.1 | Alzheimer's disease with late onset | 1% |
| I69.391 | Dysphagia following cerebral infarction | 1% |
| G80.9 | Cerebral palsy, unspecified | 1% |
| G93.40 | Encephalopathy, unspecified | 1% |
| I69.352 | Hemiplegia following cerebral infarction affecting left dominant side | 1% |
| I69.393 | Ataxia following cerebral infarction | 1% |
| I63.9 | Cerebral infarction, unspecified | 1% |
| G93.41 | Metabolic encephalopathy | 1% |
| G45.9 | Transient cerebral ischemic attack, unspecified | 1% |
| G60.9 | Hereditary and idiopathic neuropathy, unspecified | 1% |

Wounds

| ICD-10 CODE | DESCRIPTION | % OF CLINICAL GROUPING |
|-------------|--|------------------------|
| E11.621 | Type 2 diabetes mellitus with foot ulcer | 13% |
| L89.152 | Pressure ulcer of sacral region, stage 2 | 5% |
| L03.115 | Cellulitis of right lower limb | 5% |
| L89.312 | Pressure ulcer of right buttock, stage 2 | 5% |
| L89.154 | Pressure ulcer of sacral region, stage 4 | 5% |
| L03.116 | Cellulitis of left lower limb | 4% |
| L89.322 | Pressure ulcer of left buttock, stage 2 | 4% |
| L89.153 | Pressure ulcer of sacral region, stage 3 | 3% |
| T81.31XD | Disruption of external operation (surgical) wound, NEC, subs | 3% |
| S81.802D | Unspecified open wound, left lower leg, subsequent encounter | 2% |
| S81.801D | Unspecified open wound, right lower leg, subs encntr | 2% |
| E11.622 | Type 2 diabetes mellitus with other skin ulcer | 2% |
| L89.892 | Pressure ulcer of other site, stage 2 | 2% |
| S81.812D | Laceration without foreign body, left lower leg, subs encntr | 1% |
| S81.811D | Laceration w/o foreign body, right lower leg, subs encntr | 1% |
| L89.313 | Pressure ulcer of right buttock, stage 3 | 1% |
| L89.620 | Pressure ulcer of left heel, unstageable | 1% |
| L89.610 | Pressure ulcer of right heel, unstageable | 1% |
| L89.893 | Pressure ulcer of other site, stage 3 | 1% |
| L89.324 | Pressure ulcer of left buttock, stage 4 | 1% |

Complex Nursing Interventions

| ICD-10 CODE | DESCRIPTION | % OF CLINICAL GROUPING |
|-------------|---|------------------------|
| Z46.6 | Encounter for fitting and adjustment of urinary device | 55% |
| Z43.5 | Encounter for attention to cystostomy | 18% |
| Z45.2 | Encounter for adjustment and management of VAD | 9% |
| Z43.6 | Encounter for attn to oth artif openings of urinary tract | 5% |
| Z43.3 | Encounter for attention to colostomy | 5% |
| Z43.1 | Encounter for attention to gastrostomy | 3% |
| Z43.2 | Encounter for attention to ileostomy | 2% |
| Z43.0 | Encounter for attention to tracheostomy | 1% |
| Z51.11 | Encounter for antineoplastic chemotherapy | <1% |
| Z43.4 | Encounter for attn to oth artif openings of digestive tract | <1% |
| K94.09 | Other complications of colostomy | <1% |
| Z43.8 | Encounter for attention to other artificial openings | <1% |
| K94.22 | Gastrostomy infection | <1% |
| K94.19 | Other complications of enterostomy | <1% |
| T83.510D | I/I react d/t cystostomy catheter, subsequent encounter | <1% |
| K94.23 | Gastrostomy malfunction | <1% |
| K94.29 | Other complications of gastrostomy | <1% |
| K94.03 | Colostomy malfunction | <1% |
| T83.510A | I/I react d/t cystostomy catheter, initial encounter | <1% |
| K94.13 | Enterostomy malfunction | <1% |

TOP 20 DX BY CLINICAL GROUPING

Behavioral Health Care

| ICD-10 CODE | DESCRIPTION | % OF CLINICAL GROUPING |
|-------------|--|------------------------|
| F03.90 | Unspecified dementia without behavioral disturbance | 38% |
| F03.91 | Unspecified dementia with behavioral disturbance | 15% |
| F32.9 | Major depressive disorder, single episode, unspecified | 13% |
| F41.9 | Anxiety disorder, unspecified | 6% |
| F31.9 | Bipolar disorder, unspecified | 4% |
| F33.1 | Major depressive disorder, recurrent, moderate | 3% |
| F20.0 | Paranoid schizophrenia | 3% |
| F20.9 | Schizophrenia, unspecified | 2% |
| F33.9 | Major depressive disorder, recurrent, unspecified | 2% |
| F25.0 | Schizoaffective disorder, bipolar type | 2% |
| F41.1 | Generalized anxiety disorder | 1% |
| F33.2 | Major depressive disorder, recurrent severe w/o psych features | 1% |
| F25.9 | Schizoaffective disorder, unspecified | 1% |
| F33.3 | Major depressive disorder, recurrent, severe w psych symptoms | 1% |
| F41.8 | Other specified anxiety disorders | 1% |
| F29. | Unsp psychosis not due to a substance or known physiol cond | <1% |
| F25.1 | Schizoaffective disorder, depressive type | <1% |
| F33.0 | Major depressive disorder, recurrent, mild | <1% |
| F43.23 | Adjustment disorder with mixed anxiety and depressed mood | <1% |
| F43.21 | Adjustment disorder with depressed mood | <1% |

TOP 20 DX BY CLINICAL GROUPING

MMTA – Surgical Aftercare

| ICD-10 CODE | DESCRIPTION | % OF CLINICAL GROUPING |
|-------------|--|------------------------|
| Z48.812 | Encntr for surgical aftrr following surgery on the circ sys | 39% |
| Z48.3 | Aftercare following surgery for neoplasm | 24% |
| Z48.815 | Encntr for surgical aftrr following surgery on the dgstv sys | 23% |
| Z48.817 | Encntr for surgical aftrr fol surgery on the skin, subcu | 5% |
| Z48.816 | Encounter for surgical aftrr following surgery on the GU sys | 5% |
| Z48.811 | Encntr for surgical aftrr fol surgery on the nervous sys | 2% |
| Z48.813 | Encntr for surgical aftrr following surgery on the resp sys | 1% |
| Z48.22 | Encounter for aftercare following kidney transplant | <1% |
| Z48.23 | Encounter for aftercare following liver transplant | <1% |
| Z48.03 | Encounter for change or removal of drains | <1% |
| Z48.810 | Encntr for surgical aftrr fol surgery on the sense organs | <1% |
| Z48.298 | Encounter for aftercare following other organ transplant | <1% |
| Z48.21 | Encounter for aftercare following heart transplant | <1% |
| Z48.24 | Encounter for aftercare following lung transplant | <1% |
| Z48.290 | Encounter for aftercare following bone marrow transplant | <1% |
| Z48.288 | Encounter for aftercare following multiple organ transplant | <1% |
| Z48.02 | Encounter for removal of sutures | <1% |
| Z48.280 | Encounter for aftercare following heart-lung transplant | <1% |

HealthPRO® Heritage is dedicated to helping HHAs with payment reform. For more info, email homehealth@healthpro-heritage.com.
For the Top 20 Diagnoses for each Clinical Grouping for the LDS, visit healthpro-heritage.com/pdgm.

TOP 20 DX BY CLINICAL GROUPING

MMTA – Cardiac/Circulatory

| ICD-10 CODE | DESCRIPTION | % OF CLINICAL GROUPING |
|-------------|--|------------------------|
| I11.0 | Hypertensive heart disease with heart failure | 20% |
| I13.0 | Hyp hrt & chr kdny dis w hrt fail and stg 1-4/unsp chr kdny | 14% |
| I25.10 | Athscl heart disease of native coronary artery w/o ang pctrs | 9% |
| I87.2 | Venous insufficiency (chronic) (peripheral) | 8% |
| I12.9 | Hypertensive chronic kidney disease w stg 1-4/unsp chr kdny | 6% |
| I48.91 | Unspecified atrial fibrillation | 6% |
| I11.9 | Hypertensive heart disease without heart failure | 5% |
| I50.9 | Heart failure, unspecified | 3% |
| I48.0 | Paroxysmal atrial fibrillation | 3% |
| I13.2 | Hyp hrt & chr kdny dis w hrt fail and w stg 5 chr kdny/ESRD | 2% |
| I48.2 | Chronic atrial fibrillation | 2% |
| I21.4 | Non-ST elevation (NSTEMI) myocardial infarction | 2% |
| I12.0 | Hyp chr kidney disease w stage 5 chr kidney disease or ESRD | 2% |
| I73.9 | Peripheral vascular disease, unspecified | 1% |
| I26.99 | Other pulmonary embolism without acute cor pulmonale | 1% |
| I50.32 | Chronic diastolic (congestive) heart failure | 1% |
| I25.119 | Athscl heart disease of native cor art w unsp ang pctrs | 1% |
| I50.33 | Acute on chronic diastolic (congestive) heart failure | 1% |
| I50.22 | Chronic systolic (congestive) heart failure | 1% |
| I50.23 | Acute on chronic systolic (congestive) heart failure | 1% |

TOP 20 DX BY CLINICAL GROUPING

MMTA – Endocrine

| ICD-10 CODE | DESCRIPTION | % OF CLINICAL GROUPING |
|-------------|--|------------------------|
| E11.65 | Type 2 diabetes mellitus with hyperglycemia | 27% |
| E11.22 | Type 2 diabetes mellitus w diabetic chronic kidney disease | 22% |
| E11.42 | Type 2 diabetes mellitus with diabetic polyneuropathy | 15% |
| E11.40 | Type 2 diabetes mellitus with diabetic neuropathy, unsp | 15% |
| E11.51 | Type 2 diabetes w diabetic peripheral angiopath w/o gangrene | 7% |
| E11.69 | Type 2 diabetes mellitus with other specified complication | 3% |
| E11.649 | Type 2 diabetes mellitus with hypoglycemia without coma | 2% |
| E11.21 | Type 2 diabetes mellitus with diabetic nephropathy | 2% |
| E11.59 | Type 2 diabetes mellitus with oth circulatory complications | 1% |
| E10.65 | Type 1 diabetes mellitus with hyperglycemia | 1% |
| E03.9 | Hypothyroidism, unspecified | 1% |
| E11.43 | Type 2 diabetes w diabetic autonomic (poly)neuropathy | 1% |
| E11.29 | Type 2 diabetes mellitus w oth diabetic kidney complication | 1% |
| E11.610 | Type 2 diabetes mellitus w diabetic neuropathic arthropathy | 1% |
| E11.319 | Type 2 diabetes w unsp diabetic rtnop w/o macular edema | <1% |
| E10.22 | Type 1 diabetes mellitus w diabetic chronic kidney disease | <1% |
| E11.39 | Type 2 diabetes w oth diabetic ophthalmic complication | <1% |
| E11.36 | Type 2 diabetes mellitus with diabetic cataract | <1% |
| E11.00 | Type 2 diab w hyprosm w/o nonket hyprgly-hypros coma (NKHHC) | <1% |
| E11.311 | Type 2 diabetes w unsp diabetic retinopathy w macular edema | <1% |

MMTA – Gastrointestinal/Genitourinary

| ICD-10 CODE | DESCRIPTION | % OF CLINICAL GROUPING |
|-------------|---|------------------------|
| N39.0 | Urinary tract infection, site not specified | 41% |
| N40.1 | Benign prostatic hyperplasia with lower urinary tract symp | 6% |
| K92.2 | Gastrointestinal hemorrhage, unspecified | 4% |
| N31.9 | Neuromuscular dysfunction of bladder, unspecified | 4% |
| K52.9 | Noninfective gastroenteritis and colitis, unspecified | 3% |
| N30.00 | Acute cystitis without hematuria | 2% |
| K21.9 | Gastro-esophageal reflux disease without esophagitis | 2% |
| K74.60 | Unspecified cirrhosis of liver | 2% |
| N17.9 | Acute kidney failure, unspecified | 2% |
| K59.00 | Constipation, unspecified | 1% |
| K57.92 | Dvtrcli of intest, part unsp, w/o perf or abscess w/o bleed | 1% |
| K70.31 | Alcoholic cirrhosis of liver with ascites | 1% |
| K85.90 | Acute pancreatitis without necrosis or infection, unsp | 1% |
| K57.32 | Dvtrcli of lg int w/o perforation or abscess w/o bleeding | 1% |
| K81.0 | Acute cholecystitis | 1% |
| K91.2 | Postsurgical malabsorption, not elsewhere classified | 1% |
| N18.3 | Chronic kidney disease, stage 3 (moderate) | 1% |
| N10. | Acute pyelonephritis | 1% |
| K61.1 | Rectal abscess | 1% |
| N13.30 | Unspecified hydronephrosis | 1% |

MMTA – Infectious Disease/Neoplasms

| ICD-10 CODE | DESCRIPTION | % OF CLINICAL GROUPING |
|-------------|--|------------------------|
| D51.0 | Vitamin B12 defic anemia due to intrinsic factor deficiency | 7% |
| A41.9 | Sepsis, unspecified organism | 5% |
| T81.4XXD | Infection following a procedure, subsequent encounter | 5% |
| D64.9 | Anemia, unspecified | 5% |
| C61. | Malignant neoplasm of prostate | 4% |
| A04.72 | Enterocolitis due to clostridium difficile, not specified as recurrent | 3% |
| C90.00 | Multiple myeloma not having achieved remission | 3% |
| C18.9 | Malignant neoplasm of colon, unspecified | 3% |
| T81.4XXA | Infection following a procedure, initial encounter | 2% |
| C67.9 | Malignant neoplasm of bladder, unspecified | 2% |
| D50.9 | Iron deficiency anemia, unspecified | 2% |
| D51.9 | Vitamin B12 deficiency anemia, unspecified | 2% |
| C25.9 | Malignant neoplasm of pancreas, unspecified | 2% |
| C50.911 | Malignant neoplasm of unsp site of right female breast | 2% |
| D50.0 | Iron deficiency anemia secondary to blood loss (chronic) | 1% |
| C50.912 | Malignant neoplasm of unspecified site of left female breast | 1% |
| C20. | Malignant neoplasm of rectum | 1% |
| C34.92 | Malignant neoplasm of unsp part of left bronchus or lung | 1% |
| D46.9 | Myelodysplastic syndrome, unspecified | 1% |
| C15.9 | Malignant neoplasm of esophagus, unspecified | 1% |

TOP 20 DX BY CLINICAL GROUPING

MMTA – Respiratory

| ICD-10 CODE | DESCRIPTION | % OF CLINICAL GROUPING |
|-------------|--|------------------------|
| J44.9 | Chronic obstructive pulmonary disease, unspecified | 31% |
| J44.1 | Chronic obstructive pulmonary disease w (acute) exacerbation | 29% |
| J18.9 | Pneumonia, unspecified organism | 11% |
| J44.0 | Chronic obstructive pulmon disease w acute lower resp infct | 6% |
| J43.9 | Emphysema, unspecified | 3% |
| J06.9 | Acute upper respiratory infection, unspecified | 2% |
| J90. | Pleural effusion, not elsewhere classified | 2% |
| J45.909 | Unspecified asthma, uncomplicated | 1% |
| J96.21 | Acute and chronic respiratory failure with hypoxia | 1% |
| J96.11 | Chronic respiratory failure with hypoxia | 1% |
| J45.901 | Unspecified asthma with (acute) exacerbation | 1% |
| J10.1 | Flu due to oth ident influenza virus w oth resp manifest | 1% |
| J69.0 | Pneumonitis due to inhalation of food and vomit | 1% |
| J20.9 | Acute bronchitis, unspecified | 1% |
| J84.10 | Pulmonary fibrosis, unspecified | 1% |
| J18.1 | Lobar pneumonia, unspecified organism | 1% |
| J40. | Bronchitis, not specified as acute or chronic | 1% |
| J11.1 | Flu due to unidentified influenza virus w oth resp manifest | 1% |
| J96.01 | Acute respiratory failure with hypoxia | <1% |
| J15.9 | Unspecified bacterial pneumonia | <1% |

| ICD-10 CODE | DESCRIPTION | % OF CLINICAL GROUPING |
|-------------|---|------------------------|
| I10. | Essential (primary) hypertension | 53% |
| E11.9 | Type 2 diabetes mellitus without complications | 22% |
| G40.909 | Epilepsy, unsp, not intractable, without status epilepticus | 3% |
| I95.1 | Orthostatic hypotension | 2% |
| G89.29 | Other chronic pain | 2% |
| G89.4 | Chronic pain syndrome | 2% |
| T81.89XD | Oth complications of procedures, NEC, subs | 1% |
| E86.0 | Dehydration | 1% |
| T81.89XA | Oth complications of procedures, NEC, init | 1% |
| T87.89 | Other complications of amputation stump | 1% |
| E87.1 | Hypo-osmolality and hyponatremia | 1% |
| E53.8 | Deficiency of other specified B group vitamins | 1% |
| G40.89 | Other seizures | <1% |
| E78.5 | Hyperlipidemia, unspecified | <1% |
| G25.0 | Essential tremor | <1% |
| L98.8 | Oth disrd of the skin and subcutaneous tissue | <1% |
| L12.0 | Bullous pemphigoid | <1% |
| E43. | Unspecified severe protein-calorie malnutrition | <1% |
| E10.9 | Type 1 diabetes mellitus without complications | <1% |
| G40.901 | Epilepsy, unsp, not intractable, with status epilepticus | <1% |

TOP 20 DX BY CLINICAL GROUPING

Questionable Encounters

| ICD-10 CODE | DESCRIPTION | % OF CLINICAL GROUPING |
|-------------|--|------------------------|
| M62.81 | Muscle weakness (generalized) | 23% |
| R26.89 | Other abnormalities of gait and mobility | 6% |
| M54.5 | Low Back Pain | 6% |
| R26.81 | Unsteadiness on feet | 4% |
| R53.1 | Weakness | 4% |
| R29.6 | Repeated falls | 4% |
| R26.9 | Unspecified abnormalities of gait and mobility | 3% |
| M19.90 | Unspecified osteoarthritis, unspecified site | 2% |
| R26.2 | Difficulty in walking, not elsewhere classified | 2% |
| M06.9 | Rheumatoid arthritis, unspecified | 2% |
| R33.9 | Retention of urine, unspecified | 1% |
| M25.561 | Pain in right knee | 1% |
| Z91.81 | History of falling | 1% |
| Z48.89 | Encounter for other specified surgical aftercare | 1% |
| Z51.89 | Encounter for other specified aftercare | 1% |
| R55. | Syncope and collapse | 1% |
| C34.90 | Malignant neoplasm of unspecified part of unspecified bronchus or lung | 1% |
| G62.9 | Malignant neoplasm of testis | 1% |
| R42. | Dizziness and giddiness | 1% |
| M54.9 | Dorsalgia, unspecified | 1% |