



Active ingredients: prodiamine, imazaquin, simazine

# PROTECTING SOUTHERN TURF FROM COAST TO COAST



**COASTAL**  
HERBICIDE

Introducing the most all-encompassing and easy-to-use herbicide for southern turf available today. With three active ingredients combined in one jug, broad-spectrum **Coastal™ Herbicide** can be used safely on all of the four major warm-season turf grasses without the need to tank mix. This makes Coastal a smart choice when treating multiple lawns. And Coastal outperforms the current industry leader on hard-to-control weeds, including crabgrass and *Poa annua*.



## COASTAL IS THE ULTIMATE HERBICIDE FOR WARM-SEASON TURF GRASSES

- » Broad-spectrum, pre- and post-emergent broadleaf and grassy weed control
  - » For use on golf courses, residential and commercial turf, athletic fields and sod farms
  - » Safe to use on Bermudagrass, centipede grass, St. Augustine grass and Zoysia grass
  - » Three active ingredients/modes of action broaden weed control spectrum and reduce mixing time, potential mistakes and jug disposal
  - » Excellent resistance management tool
  - » Low lateral mobility
  - » Most economical *Poa annua* control available
- 

# A TOP PERFORMER IN UNIVERSITY TRIALS ACROSS THE SOUTH

In preliminary trials conducted by Mississippi State University, University of Georgia, University of Tennessee, North Carolina State University, University of Florida, Louisiana State University, and Texas A&M University, Coastal Herbicide was shown to provide exceptional control of *Poa annua* and crabgrass, in addition to control of broadleaf weeds like white clover, henbit, lawn burweed and others.

## WEED CONTROL COMPARISON

COASTAL CONTROL	Pre	-	-	-	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	
	Post	S	S	S	-	-	S	C	S	S	C	C	S	S	C	C	S	C	C	
SPECTICLE CONTROL	Pre	-	C	C	C	C	C	C	C	C	-	-	-	C	C	C	C	-	C	C
	Post	-	S	-	S	-	S	S	-	-	-	-	-	S	-	-	C	-	-	-
	Weed Type	Perennial Sedges	Kyllinga	Annual Sedges	Crabgrass	Goose Grass	Rye Grass	Poa	Lawn Burweed	Field Sandbur	Purple Deadnettle	Wild Garlic/Onions	Carpetweed	Chickweed	White Clover	Corn Speedwell	Doveweed	Dollarweed	Henbit	Spurge

**C** indicates control at labeled use rates

**S** indicates suppression or control on newly emerged plants up to the 1-2 leaf stage at the high use rate

**■** indicates advantage

**■** indicates disadvantage

## TIMING AND USE RATES

The use rates of this product range from 48 oz. per acre up to the maximum rate of 64 oz. per acre. The length of control, size and spectrum of weeds controlled post-emergence increase with higher use rates. The use of a spray adjuvant is required to obtain post-emergence weed control. Mix recommended rate of this product with water and apply in a spray volume of 10 to 40 gallons per acre.

**FALL AND WINTER** | 64 oz. per acre. Do not apply before September 15.

**JANUARY THROUGH MARCH** | 48 oz. per acre. Higher rates may be used if spring transition is at least 4 to 6 weeks away. Do not apply if spring transition is imminent or expected within 3-4 weeks unless delayed green-up and/or temporary yellowing can be tolerated.

**MAY** | Applications that occur after full spring transition may use any labeled rate. Do not use after May 31.

For more information about Coastal, talk to your Sipcarn Agro distributor or visit [coastalherbicide.com](http://coastalherbicide.com).

