

Spinnaker

ACTIVE INGREDIENT: COPPER HYDROXIDE

PERFORMANCE HIGHLIGHTS

- Spinnaker provides superior disease control with lower metallic use rates
- Proven control of a wide range of bacterial diseases and fungal pathogens
- Outstanding dry flowable formulation for easy handling, mixing and use along with most insecticides, fungicides and foliar nutrients
- Superior coverage and rainfastness
- Low risk for resistance

Chemical Class: Copper Hydroxide

Packaging: 25 lbs bags

Pre-Harvest Interval: 0 days



KEY CROPS

Sugarbeets

Potatoes

KEY DISEASES CONTROLLED

Cercospora leaf spot

Early Blight

Late Blight

SUGARBEET AND POTATOES FIELD CROPS

Crop	Disease	Application Rate/Acre	Use Instructions
Potato	Early Blight Late Blight	0.5-1.75 lb	Apply 0.5 to 1.75 pounds at 5- to 10-day intervals if needed starting when plants are 2 to 6 inches high in locations where disease is light. Apply up to 1.75 pounds per acre when is more severe. Under conditions of severe disease, control with Spinnaker will be improved by tank mixing with other compatible fungicides registered for use on potatoes. Read and follow all label instructions of tank mix partners.
Sugar Beet	Cercospora Leaf Spot	0.75 - 2.0 lb	Begin applications when conditions first favor disease development and repeat at 10- to 14-day intervals if needed. Use the higher rates when conditions favor disease. Add a spreader/sticker. NDSU recommend tank mixing Spinnaker with other compatible fungicides registered for use on sugar beets, such as Minerva or Miramar.



sipcamagro.com

1-877-898-9514

© 2019 Sipcam Agro USA, Inc. Always read and follow label directions and precautions for use. Product(s) referenced herein may not be registered in all states. Confirm your state's registration with your Sipcam Agro USA, Inc. representative.



SIPCAM AGRO
USA, INC.