



**TECHNICAL INFORMATION BULLETIN**  
**FOR THE USE OF STARGUS® BIOFUNGICIDE**  
 Bacterial spot (*Xanthomonas campestris pv. pruni*) on stone fruit  
 Fire blight (*Erwinia amylovora*) on pome fruit  
 Powdery mildew (*Podosphaera leucotricha*)

**FIFRA Section 2(ee) Recommendation**

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U. S. Environmental Protection Agency.

FOR USE IN ALL STATES WHERE CROP AND PEST(S) EXIST AND WHERE 2(EE) RECOMMENDATIONS ARE RECOGNIZED.

**STARGUS®**

**EPA Reg. No. 84059-28**

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Crop	Target Disease	Application Method	Product Use Rate per Application	Application Instructions
Stone Fruit	Bacterial blight ( <i>Xanthomonas campestris pv. pruni</i> )	Foliar (Ground) Applications from shuck split to final cover spray	2 quarts per acre	Applications beginning at shuck split should be made in a minimum of 75 gallons of water per acre using spray equipment designed to achieve full coverage of orchard crops. Water volume should not exceed 250 gallons of water per acre. The use of a non-penetrating spreader-sticker may be used to improve coverage and rain fastness. Spray interval between applications should not exceed 10 days.
Pome Fruit	Fire blight ( <i>Erwinia amylovora</i> )  Powdery mildew <i>Podosphaera leucotricha</i>	Foliar (Ground) Applications	2-4 quarts per acre	Fire Blight – Apply this product at 2 – 4 quarts per acre in 50 - 100 gallons of water per acre beginning at green tip and continuing through petal fall. For maximum control, use this product prior to infection events. During periods of rapid development and frequent infection periods, use spray intervals of 3 - 7 days.  Apply in sufficient water to provide full coverage. For improved performance, use this product in a rotational program with antibiotics registered for fire blight control such as but not limited to oxytetracycline or streptomycin. Use caution when selecting spray adjuvants. Select only those adjuvants which through prior experience do not affect



				fruit finish when combined with this product
STARGUS® has a <b>pre-harvest interval (PHI)</b> of <b>0 days</b> . Do not enter or allow worker entry into treated areas during the <b>restricted-entry interval (REI)</b> of <b>4 hours</b> .				

**All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.**

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and carefully observe the cautionary statements plus all the other information appearing on the product label.

**This recommendation must be in the possession of the user at the time of pesticide application.**

STARGUS® is a registered trademark of Marrone Bio Innovations, Inc.

© Copyright Marrone Bio Innovations, Inc., 2018

Marrone Bio Innovations, Inc.

1540 Drew Ave.

Davis, California 95618

Telephone (530) 750-2800

[www.marronebioinnovations.com](http://www.marronebioinnovations.com)

Version May 5, 2018