

**NUGOLD®**  
**FLORFENICOL 30% INJECTION**  
**VETERINARY USE**

**COMPOSITION**

Eac 1 ml contains:

Florfenicol	<b>300 mg</b>
Polietilenglicol-200 c.s.p.	1mL

**INDICATIONS** In cattle and pigs for the treatment of infections caused by:

**Cattle:** *Mannheimia haemolytica*, *Pasteurella multocida*, *Fusobacterium necrophorum* and *Bacteroides melaninogenicus*.

**Swines:** *Pasteurella multocida*, *Actinobacillus pleuropneumoniae* and *mycoplasma hyopneumoniae*.

**DOSAGE AND ADMINISTRATION**

**Bovines**

Administer intramuscular route (preferably in the neck table) 1mL per 15 Kg of weight, which corresponds to the recommended dose of Florfenicol, 20 mg per kg of weight, repeat the dose at 48 hours. Don't administer more than 10 mL per application site. Alternatively, 2 mL can be administered subcutaneously for every 15 Kg of weight (40mg / kg) as a single dose, seeking Long-acting effect.

**Pigs**

Administer 1mL deep intramuscular route for every 20 Kg of weight, which corresponds to the recommended dose of 15mg of Florfenicol per kg of weight, repeat the dose at 48 hours. Do not administer more than 3mL per application site.

**WITHDRAWAL TIME**

Cattle and pigs treated intramuscularly should not be slaughtered for human consumption until 30 days after the end of treatment. Cattle treated subcutaneously should not be slaughtered for human consumption until 44 days after the end of treatment. It is not allowed to be used in cattle producing milk for human consumption.

**RECOMMENDATIONS**

It's recommended to treat animals in the early stages of the disease and evaluate the response to treatment within 48 hours after the second application. If the clinical signs of respiratory disease persist 48 hours after the last application, the antibiotic should be changed and treated until the signs disappear.

### **PRECAUTIONS**

Veterinary use.

Keep out of reach of children.

Store in a cool dry place, protected from light.

Sale under the formula of the veterinarian.

### **CONTRAINDICATIONS**

Do not use in pregnant or lactating females.

Do not use in animals with renal insufficiency.

Do not use during vaccination periods or in animals with cases of immunosuppression.

**COMMERCIAL PRESENTATIONS:** Vial bottle for 20, 50, 100, 250 and 500mL.

**ICA REGISTRY No.: 9682-MV**

Registered trademark of  
Rotam Agrochem International Co. Ltd.