










## Routine Immunization Schedule

| Age  | Vaccine(s) your child / you will get   | Diseases your child / you will be protected against   |
|--|--|---|
|  <b>2 months</b>  | <ul style="list-style-type: none"> <li>DTaP-IPV-Hib-HB*</li> <li>Pneumococcal conjugate (PNEU-C13)</li> <li>Rotavirus</li> </ul>   | <ul style="list-style-type: none"> <li>Diphtheria, tetanus, pertussis (whooping cough), polio, <i>Haemophilus influenzae</i> type b, hepatitis B</li> <li>Pneumococcal Disease</li> <li>Rotavirus</li> </ul>                                |
|  <b>4 months</b>  | <ul style="list-style-type: none"> <li>DTaP-IPV-Hib-HB*</li> <li>Pneumococcal conjugate (PNEU-C13)</li> <li>Meningococcal conjugate (MenconC)</li> <li>Rotavirus</li> </ul>  | <ul style="list-style-type: none"> <li>Diphtheria, tetanus, pertussis (whooping cough), polio, <i>Haemophilus influenzae</i> type b, hepatitis B</li> <li>Pneumococcal Disease</li> <li>Meningococcal Disease</li> <li>Rotavirus</li> </ul> |
|  <b>6 months</b>  | <ul style="list-style-type: none"> <li>DTaP-IPV-Hib-HB*</li> <li>Pneumococcal conjugate (PNEU-C13) (for high-risk children only)</li> <li>Rotavirus</li> </ul>   | <ul style="list-style-type: none"> <li>Diphtheria, tetanus, pertussis (whooping cough), polio, <i>Haemophilus influenzae</i> type b, hepatitis B</li> <li>Pneumococcal Disease</li> <li>Rotavirus</li> </ul>                                |
|  <b>12 months</b> | <ul style="list-style-type: none"> <li>MMR-Var</li> <li>Pneumococcal conjugate (PNEU-C13)</li> <li>Meningococcal conjugate (MenconC)</li> </ul>  | <ul style="list-style-type: none"> <li>Measles, mumps, rubella, varicella (chickenpox)</li> <li>Pneumococcal Disease</li> <li>Meningococcal Disease</li> </ul>  |
|  <b>18 months</b> | <ul style="list-style-type: none"> <li>DTaP-IPV-Hib</li> </ul>   | <ul style="list-style-type: none"> <li>Diphtheria, tetanus, pertussis (whooping cough), polio, <i>Haemophilus influenzae</i> type b</li> </ul>  |
|  <b>4 years</b> | <ul style="list-style-type: none"> <li>dTap-IPV</li> <li>MMR-Var</li> </ul>  | <ul style="list-style-type: none"> <li>Diphtheria, tetanus, pertussis (whooping cough), polio</li> <li>Measles, mumps, rubella, varicella (chickenpox)</li> </ul>   |
|  <b>Grade 6</b> | <ul style="list-style-type: none"> <li>Hepatitis B (3 doses over 6 months)</li> <li>HPV (2 or 3 doses over 6 months)</li> </ul>  | <ul style="list-style-type: none"> <li>Hepatitis B</li> <li>Human papillomavirus</li> </ul>   |
|  <b>Grade 9</b> | <ul style="list-style-type: none"> <li>dTap</li> <li>MenC-ACYW</li> </ul>  | <ul style="list-style-type: none"> <li>Diphtheria, tetanus, pertussis (whooping cough)</li> <li>Meningococcal Disease</li> </ul>  |
|  <b>Adult</b>   | <ul style="list-style-type: none"> <li>Booster doses: dTap (1 dose) then Td every 10 years</li> <li>Hepatitis B (3 doses for unprotected adults born in 1981 or later)</li> <li>MMR (1 or 2 doses if born in 1957 or later and no record showing given in childhood)</li> <li>Varicella (chickenpox) for unprotected adults (2 doses)</li> <li>Pneumococcal polysaccharide (PNEUMO-P) at 65 years of age or older</li> </ul> | <ul style="list-style-type: none"> <li>Diphtheria, tetanus, pertussis (whooping cough)</li> <li>Hepatitis B</li> <li>Measles, mumps, rubella</li> <li>Varicella (chickenpox)</li> <li>Pneumococcal Disease</li> </ul>                       |
| <b>Everyone (6 months of age and older)</b>  | <ul style="list-style-type: none"> <li>Annual (seasonal) influenza</li> </ul>  | <ul style="list-style-type: none"> <li>Influenza</li> </ul>   |

For more information, go to the Adult Immunization information sheet on [immunizealberta.ca](http://immunizealberta.ca)

\* DTaP-IPV-Hib-HB – for children born March 1, 2018 or later. Children born before March 1, 2018 will get DTaP-IPV-Hib at 2, 4 and 6 months and Hepatitis B in school.

**Note:** More vaccines may be needed because of health conditions, work, school, living arrangements, household contacts, lifestyle risks, travel, or because a person was not fully immunized in childhood. Check with your healthcare provider, a public health nurse, your workplace health and safety department, or a travel health clinic (e.g., [AHS Travel Health Services](http://AHS Travel Health Services)) to find out if you need any other vaccines and if they are free for you.