

**SUGGESTED FORMULA**

Baclofen 30mg / Lidocaine 50mg Suppositories

Version number: 1.0

Volume or quantity: #10 Suppositories

Baclofen, USP	0.3gm
Lidocaine, USP	0.5gm
Witepsol [®] H-15 Base	19.2gm

*Formulation is based on a mold size that holds 2gm, it is important to calibrate the molds being used and adjust formula accordingly before compounding.

SUGGESTED COMPOUNDING PROCEDURES

Weigh all ingredients

1. Weight Witepsol and gently heat in a beaker
2. Once base is melted add Baclofen and Lidocaine
3. Once mixture is homogenous pour the mixture into suppository molds and allow to cool
4. Suggested Quality assessments
 - a. % variability & % deviation from theoretical <10% - (remake if >10%)
 - b. Surface appearance and feel
 - c. Melting test
 - d. Quantity
 - e. Label - auxiliary labels, storage, BUD, compounded medication

No claims are made as to the safety or efficacy of this preparation. This formulation is provided solely at the unsolicited request of the pharmacist.

Beyond-Use Dates of preparations are conservative estimates from reference books, peer-reviewed literature, and intended duration of therapy, formulation from commercially available products, organoleptic observations and current USP guidelines. Compounders may have stability studies performed by a reputable laboratory if they wish to extend the Beyond-Use Date. It is recommended that you follow USP <795> recommendations for potency testing.

Beyond-Use Date should be based on current USP <795> guidelines

Precautions:

Precautions should be taken to prevent cross-contamination and exposure of ingredients to the compounder and contamination of the preparation by the compounder. Wear appropriate protective equipment. Use safety enclosures (hoods) when weighing and mixing

4/18JD