

**SUGGESTED FORMULA**

Chlorpromazine Hydrochloride 25mg in Polyethylene Glycol (1000/3350) Suppositories

Version number: 1.0

Volume or quantity: #100 Suppositories

Chlorpromazine Hydrochloride	2.5gm
Polyethylene Glycol 1000, NF (PO118)*	75gm
Polyethylene Glycol 3350, NF (PO125)*	25gm

SUGGESTED COMPOUNDING PROCEDURES

*It is important to calibrate the suppositories molds you are using and adjust formulation accordingly.

Weigh all ingredients

1. Calculate the required quantity of each ingredient for the total amount to be prepared
2. Accurately weigh and/or measure each ingredient
3. Melt the Polyethylene Glycol's to about 55°C mixing well
4. Triturate the Chlorpromazine Hydrochloride to a fine powder with a mortar and pestle
5. Sprinkle the powder on the melted base and stir until thoroughly mixed
6. Pour into previously calibrated molds and allow to cool
7. Package and Label
8. Suggested Quality assessments
 - a. % variability & % deviation from theoretical <10% - (remake if >10%)
 - b. Surface appearance and feel
 - c. Melting test
 - d. Quantity
 - e. Label - auxiliary labels, storage, BUD, compounded medication

Package in air tight, Light resistant containers and Store Refrigerated

No claims are made as to the safety or efficacy of this preparation. This formulation is provided solely at the unsolicited request of the pharmacist.

Beyond-Use Dates of preparations are conservative estimates from reference books, peer-reviewed literature, and intended duration of therapy, formulation from commercially available products, organoleptic observations and current USP guidelines. Compounders may have stability studies performed by a reputable laboratory if they wish to extend the Beyond-Use Date.

It is recommended that you follow USP <795> recommendations for potency testing.

Beyond-Use Date estimated to be **6 Months per CompoundingToday.com**

Precautions:

Precautions should be taken to prevent cross-contamination and exposure of ingredients to the compounder and contamination of the preparation by the compounder. Wear appropriate protective equipment. Use safety enclosures (hoods) when weighing and mixing

9/18 JD