



SUGGESTED FORMULA

JESSNER'S TOPICAL SOLUTION

Version number: 1.0

Volume: 100 mL

Salicylic Acid, USP Crystalline Powder	14 g
Resorcinol USP	14 g
Lactic Acid USP 88%	14 mL
Ethyl Alcohol, 190 Proof USP	Q.S. 100 mL

SUGGESTED COMPOUNDING PROCEDURES

1. Add approximately 65% of the final volume of Ethyl Alcohol to appropriate beaker with a stir bar.
2. Add Salicylic Acid and Resorcinol to beaker and stir to dissolve.
3. Add Lactic Acid to step 2 and mix.
4. Bring to final volume with Ethyl Alcohol and mix until uniform
5. Suggested quality assessments:
 - a. Clarity
 - b. Color
 - c. Label (s)- auxiliary labels (external use), storage, BUD

Store in air-tight container.

Store at Controlled Room Temperature

No claims are made as to the safety or efficacy of this preparation. This formulation is provided solely at the unsolicited request of the pharmacist.

Beyond-Use Dates of preparations are conservative estimates by the formulator using reference books, peer-reviewed literature, intended duration of therapy, formulation from commercially available products, organoleptic observations and current USP guidelines. Compounders may have stability studies performed by a reputable laboratory if they wish to extend the Beyond-Use Date.

It is recommended that you follow USP <795> recommendations for potency testing.

Beyond-Use Date is estimated to be 30 days.

Precautions should be taken to prevent cross-contamination and exposure of ingredients to the compounder and contamination of the preparation by the compounder. Wear appropriate protective equipment . Use safety enclosures (hoods) when weighing and mixing.

05/15rd