



---

### SUGGESTED FORMULA

---

#### Methimazole 10mg/mL Suspension – Oil Formula

Version number: 1.0

Volume: 50 mL

---

Methimazole, USP (M1137)	0.5gm
Flavor, Powder	0.5 gm, OR
Flavor, Oil Soluble	2.5 mL
Corn Oil, NF (CO136)	50mL Q.S.

---

\*If using a powder flavor, add powder to step#2 and dry triturate with Methimazole. If using a liquid flavor add directly to the bottle

#### SUGGESTED COMPOUNDING PROCEDURES

1. Weigh Methimazole and place in glass mortar and pestle
2. Dry triturate to break down any clumps in the chemical
3. Add enough oil into mortar and pestle to form a “paste”
4. Triturate until all Methimazole is broken down and incorporated into liquid
5. Add more oil until pourable mixture and pour into calibrated final container
6. Perform liquid “washes” in mortar and pestle to integrate remaining chemical
7. Pour each “wash” into final container taking precaution not to go over the final volume
8. Q.S. medication and shake well.
8. Suggested Quality Assessments – follow pharmacy SOPs:
  - a. Weight to Volume calculation
  - b. Color
  - c. Pourability
  - d. Settling
  - e. Resuspendability

Store in air tight amber plastic containers

Store Room Temperature

No claims are made as to the safety or efficacy of this preparation. This formulation is provided solely at the unsolicited request of the pharmacist.

Beyond-Use Dates of preparations are conservative estimates by the formulator using reference books, peer-reviewed literature, and intended duration of therapy, formulation from commercially available products, organoleptic observations and current USP guidelines. Compounders may have stability studies performed by a reputable laboratory if they wish to extend the Beyond-Use Date. It is recommended that you follow USP <795> recommendations for potency testing.

**Beyond-Use Date is estimated to be 180 days** Per USP <795> for non-aqueous formulas - consult current version of USP General Chapter <795>.

Precautions should be taken to prevent cross-contamination and exposure of ingredients to the compounder and contamination of the preparation by the compounder. Wear appropriate protective equipment. Use safety enclosures (hoods) when weighing and mixing.

10/18 JD