



SUGGESTED FORMULA

Methocarbamol 375mg/mL in Ora-Plus™ and Ora-Sweet™

Version number: 1.0

Volume: 100 mL

Methocarbamol, USP (M1088)	37.5gm
Vehicle for Oral Suspension, NF (Ora-Plus™)	38mL
Peppermint Oil, NF (PE105)	2mL
Vehicle for Oral Solution, NF (Ora-Sweet™)	q.s. 100mL

***See formulation for Oral Suspension and Oral Solution on SpectrumRx.com under Formulations**

SUGGESTED COMPOUNDING PROCEDURES

1. Calculate the required quantity of each ingredient for the total amount to be prepared
2. Accurately weigh and/or measure each ingredient
3. Add Methocarbamol and add to mortar and pestle
4. Dry triturate to break down any clumps in the material
5. Add Ora-Plus™ to final container
6. Add enough Ora-Sweet™ into mortar and pestle to form a “paste”
7. Triturate until all Methocarbamol is broken down and incorporated into liquid
8. Add more Ora-Sweet™ until pourable mixture and pour into Step 5 (calibrated final container)
9. Add flavor to bottle
10. Perform liquid “washes” in mortar and pestle to integrate remaining chemical
11. Pour each “wash” into final container taking precaution not to go over the final volume
12. Bring to final volume with Ora-Sweet™
13. Package and label
 9. Suggested Quality Assessments – follow pharmacy SOPs:
 - a. Weight to Volume calculation
 - b. Pourability
 - c. Settling/Resuspendability
 - d. Label - auxiliary labels, storage, BUD, compounded medication

Store in air tight amber plastic containers; Refrigerate

No claims are made as to the safety or efficacy of this preparation. This formulation is provided solely at the unsolicited request of the pharmacist.

Beyond-Use Dates of preparations are conservative estimates by the formulator using reference books, peer-reviewed literature, and intended duration of therapy, formulation from commercially available products, organoleptic observations and current USP guidelines. Compounders may have stability studies performed by a reputable laboratory if they wish to extend the Beyond-Use Date. It is recommended that you follow USP <795> recommendations for potency testing.

Beyond-Use Date of 14 Days can be used for this preparation when stored in a refrigerator

Precautions should be taken to prevent cross-contamination and exposure of ingredients to the compounder and contamination of the preparation by the compounder. Wear appropriate protective equipment. Use safety enclosures (hoods) when weighing and mixing.

7/18 JD