



SUGGESTED FORMULA

Neuropathic Intra-Oral AG/ORA Pregab and Ephedrine (4/1/4/4/4/10/0.2 %

Version number: 1.0

Volume: 30gm

Carbamazapine ,USP (CA250)	1.2GM
Lidocaine Hydrochloride,USP(LI103)	0.3GM
Ketamine Hydrochloride,USP(K1068)	1.2GM
Ketoprofen,USP(K1155)	1.2GM
Gabapentin, USP(G1092)	1.2 GM
Lyrica [®] 300mg Capsules (Commercial)	10 Capsules
Ephedrine Sulfate, USP(EP120)	0.06GM
Kris Anhydrous Gel	15 GM
Oral Paste	30 GM Q.S.

SUGGESTED COMPOUNDING PROCEDURES

1. Triturate all powders together
2. Open Lyrica[®] capsules and add to Step 1
3. Mix Kris Anhydrous Gel into Step 2
4. Q.S. with orabase and mix well
5. Mill all ingredients
6. Check for “grit” by rubbing a small amount on the back of your gloved hand, mill again if necessary

Suggested Quality Assurance Checklist:

% Calculation: _____

No Grit: _____

Correct Auxiliary Labels: _____

Room Temperature

No claims are made as to the safety or efficacy of this preparation. This formulation is provided solely at the unsolicited request of the pharmacist.

Beyond-Use Dates of preparations are conservative estimates by the formulator using reference books, peer-reviewed literature, and intended duration of therapy, formulation from commercially available products, organoleptic observations and current USP guidelines. Compounders may have stability studies performed by a reputable laboratory if they wish to extend the Beyond-Use Date.

It is recommended that you follow USP <795> recommendations for potency testing.

Beyond-Use Date is estimated to be 30 days

Hazardous drugs should be handled according to USP <800>, NIOSH standards and pharmacy SOPs. This preparation should be compounded in a biological safety cabinet or compounding aseptic containment isolator. Precautions should be taken to prevent cross-contamination and exposure of ingredients to the compounder and contamination of the preparation by the compounder. Wear appropriate personal protective equipment.

05/18 JD