



SUGGESTED FORMULA

Neuropathic Intra-oral AG Pregab (4/1/4/4/4/10) %

Version number: 1.0

Volume: 30gm

Carbamazapine ,USP	1.2 GM
Lidocaine Hydrochloride,USP	0.3 GM
Ketamine Hydrochloride,USP	1.2 GM
Ketoprofen,USP	1.2 GM
Gabapentin, USP	1.2 GM
Lyrica [®] 300mg Capsules(Commercial)	10 Capsules
Kris Anhydrous Gel	30 GM Q.S.

SUGGESTED COMPOUNDING PROCEDURES

1. Triturate all powders together
2. Open Lyrica[®] capsules and add to Step1
2. Mix Kris Anhydrous Gel into Step2
3. Mill all ingredients
4. Check for “grit” by rubbing a small amount on the back of your gloved hand, mill again if necessary

Suggested Quality Assurance Checklist:

% Calculation: _____

No Grit: _____

Correct Auxiliary Labels: _____

Room Temperature

No claims are made as to the safety or efficacy of this preparation. This formulation is provided solely at the unsolicited request of the pharmacist.

Beyond-Use Dates of preparations are conservative estimates by the formulator using reference books, peer-reviewed literature, intended duration of therapy, formulation from commercially available products, organoleptic observations and current USP guidelines. Compounders may have stability studies performed by a reputable laboratory if they wish to extend the Beyond-Use Date.

It is recommended that you follow USP <795> recommendations for potency testing.

Beyond-Use Date is estimated to be 180 days

Precautions should be taken when with hazardous drugs. These may be absorbed through skin, mucous membranes and lungs. Wear appropriate personal protective equipment – and inform personnel of the risk. Mix and prepare in vented enclosure (hood), USP standards and facility SOPs.

06/15 JD