



SUGGESTED FORMULA

Nicotine Polacrilex 2mg Lollipops

Version number: 1.0

Volume: #1 lollipop

Nicotine Polacrilex	1.1gm
Ethyl Alcohol 95%, USP	17mL
Stevoside, NF	1.25gm
Flavor of choice (Optional)	0.5mL
Sorbitol Lollipop Base	Q.S.

Note: Nicotine Polacrilex 1.1gm is equivalent to 200mg of Nicotine. The size of the lollipop mold must be determined before the final weight of the base that will be used is selected.

SUGGESTED COMPOUNDING PROCEDURES

1. Calculate the required quantity of each ingredient for the total amount to be prepared
2. Accurately weigh and/or measure each ingredient
3. Disperse the nicotine polacrilex in the alcohol
4. Melt the Sorbitol candies at a temperature that is no higher the 70°C
5. Add the nicotine dispersion and mix well
6. Cool the mixture slightly and pour it into lollipop molds while the mixture is still fluid
7. Allow the mixture to cool, and remove it from the molds. The lollipops can be dusted with powdered sugar if they are sticky
8. Suggested quality assessments:
 - a. Appearance, feel and melting test
 - b. Color
 - c. Label (s)- auxiliary labels (Keep out of the reach of children), storage, BUD

Store in light resistant air-tight container.

Store at Controlled Room Temperature

No claims are made as to the safety or efficacy of this preparation. This formulation is provided solely at the unsolicited request of the pharmacist.

Beyond-Use Dates of preparations are conservative estimates by the formulator using reference books, peer-reviewed literature, intended duration of therapy, formulation from commercially available products, organoleptic observations and current USP guidelines. Compounders may have stability studies performed by a reputable laboratory if they wish to extend the Beyond-Use Date.

It is recommended that you follow USP <795> recommendations for potency testing.

Beyond-Use Date is estimated to be 6months per Compounding Today.com

Precautions should be taken to prevent cross-contamination and exposure of ingredients to the compounder and contamination of the preparation by the compounder. Wear appropriate protective equipment. Use safety enclosures (hoods) when weighing and mixing.

9/15 JD