



SUGGESTED FORMULA

Pluronic[™] Lecithin Organogel Vehicle

Version number: 1.0

Volume: 100 mL

Lecithin Isopropyl Palmitate Solution	20 mL
Pluronic [™] F-127 20% Gel	80 mL

SUGGESTED COMPOUNDING PROCEDURES

1. Accurately measure each ingredient
2. Mix the two viscous liquids together well, in beaker, between two syringes or in a mortar with pestle.
3. Mixture will solidify as it warms to room temperature
 - a. Weight to Volume calculation
 - b. Uniformity
 - c. Texture

Store Refrigerated or at room temperature.

No claims are made as to the safety or efficacy of this preparation. This formulation is provided solely at the unsolicited request of the pharmacist.

Beyond-Use Dates of preparations are conservative estimates by the formulator using reference books, peer-reviewed literature, intended duration of therapy, formulation from commercially available products, organoleptic observations and current USP guidelines. Compounders may have stability studies performed by a reputable laboratory if they wish to extend the Beyond-Use Date. It is recommended that you follow USP <795> recommendations for potency testing.

A beyond-use date of 30 days can be applied to this preparation, per www.compoundingtoday.com

Precautions should be taken to prevent cross-contamination and exposure of ingredients to the compounder and contamination of the preparation by the compounder. Wear appropriate protective equipment. Use safety enclosures (hoods) when weighing and mixing.

01/16 RD