



SUGGESTED FORMULA

Podophyllum Resin 1% and Salicylic Acid 3% in Flexible Collodion Suspension

Version number: 1.0

Volume: 100mL

Podophyllum Resin, USP (PO105)	1gm
Salicylic Acid, USP (SA130)	3gm
Dehydrated Alcohol, 200 Proof, Undenatured USP (ET107)	10mL
Flexible Collodion, USP (CO125)	Q.S. 100mL

SUGGESTED COMPOUNDING PROCEDURES

1. Calculate the required quantity of each ingredient for the total amount to be prepared
2. Accurately weigh and/or measure each ingredient
3. Make paste with the Podophyllin Resin and 5mL Dehydrated Alcohol
4. Dissolve the Salicylic Acid in 5mL Dehydrated Alcohol
5. Geometrically incorporate the Podophyllin paste into 80mL Flexible Collodion
6. Add sufficient Flexible Collodion to make 100mL total volume
7. Package and label
8. Suggested Quality Assessments – follow pharmacy SOPs:
 - a. Weight to Volume calculation
 - b. Color
 - c. Pourability
 - d. Settling
 - e. Resuspendability

Store in air tight amber plastic containers at room temperature

No claims are made as to the safety or efficacy of this preparation. This formulation is provided solely at the unsolicited request of the pharmacist.

Beyond-Use Dates of preparations are conservative estimates by the formulator using reference books, peer-reviewed literature, and intended duration of therapy, formulation from commercially available products, organoleptic observations and current USP guidelines. Compounders may have stability studies performed by a reputable laboratory if they wish to extend the Beyond-Use Date. It is recommended that you follow USP <795> recommendations for potency testing.

Beyond-Use Date is estimated to be 180 Days Per *CompoundingToday.com* Formula #887

Precautions should be taken to prevent cross-contamination and exposure of ingredients to the compounder and contamination of the preparation by the compounder. Wear appropriate protective equipment. Use safety enclosures (hoods) when weighing and mixing.

6/18 JD