



Salicylic Acid 30% Topical Peel

Version number: 1.0

Volume: 30 g

Salicylic Acid, USP Crystalline Powder	9 g
Polysorbate 60 NF	2 g
Ethyl Alcohol, 200 Proof USP	19 g

SUGGESTED COMPOUNDING PROCEDURES

1. In a beaker mix Salicylic Acid, Ethyl Alcohol and Polysorbate 60 with stir bar.
(the specific gravity of Polysorbate 60 is 1.10 at 25°C, therefore 1 g is equivalent to 0.91 mL).
2. Cover beaker with aluminum foil and stir until all ingredients have completely dissolved.
3. Suggested quality assessments:
 - a. Clarity
 - b. Color
 - c. Label (s)- auxiliary labels (external use), storage, BUD

Protect from light.

Store in air-tight container.

Store at Controlled Room Temperature

No claims are made as to the safety or efficacy of this preparation. This formulation is provided solely at the unsolicited request of the pharmacist.

Beyond-Use Dates of preparations are conservative estimates by the formulator using reference books, peer-reviewed literature, intended duration of therapy, formulation from commercially available products, organoleptic observations and current USP guidelines. Compounders may have stability studies performed by a reputable laboratory if they wish to extend the Beyond-Use Date.

It is recommended that you follow USP <795> recommendations for potency testing.

According to USP guidelines, the storage period for water containing formulations is 14 days when stored between 2° and 8 °C. See USP Chapter <795>

Beyond-Use Date is estimated to be 14 days, stored refrigerated.

Precautions should be taken to prevent cross-contamination and exposure of ingredients to the compounder and contamination of the preparation by the compounder. Wear appropriate protective equipment . Use safety enclosures (hoods) when weighing and mixing.

05/15rd