



### SUGGESTED FORMULA

---

#### Sorbitol 70% Lollipop Vehicle Base

Version number: 1.0

Quantity: 100mL

---

Sorbitol Solution 70% Solution, USP (SO220)	99mL
Saccharin Sodium (Anhydrous)	0.150gm
Active Drug	X
Flavor of Choice	X

---

### SUGGESTED COMPOUNDING PROCEDURES

1. Calculate the required quantity of each ingredient for the total amount to be prepared
2. Accurately weight and/or measure each ingredient
3. Heat the Sorbitol Solution slowly until it reaches 150°C
4. Remove from the heat. When the temperature decreases to approximately 100°C, incorporate the active drug, Saccharin sodium and flavor
5. Pour into lollipop molds that have been previously spray with a vegetable spray such as PAM™ providing a thin coating
6. Allow to cool and preparation to set
7. Package and label

Store in air-tight container, at Room Temperature

No claims are made as to the safety or efficacy of this preparation. This formulation is provided solely at the unsolicited request of the pharmacist.

Beyond-Use Dates of preparations are conservative estimates from reference books, peer-reviewed literature, and intended duration of therapy, formulation from commercially available products, organoleptic observations and current USP guidelines. Compounders may have stability studies performed by a reputable laboratory if they wish to extend the Beyond-Use Date.

It is recommended that you follow USP <795> recommendations for potency testing.

Beyond-Use Date estimated to be **6 Months** per CompoundingToday.com Formula#1670

**General:** Precautions should be taken to prevent cross-contamination and exposure of ingredients to the compounder and contamination of the preparation by the compounder. Wear appropriate protective equipment. Use safety enclosures (hoods) when weighing and mixing.

6/18JD