



SUGGESTED FORMULA

Triamcinolone 0.1%/ Ketoconazole 1%/ Gentamicin 0.3% in Lanolin

Version number: 1.0

Volume or quantity: 60 gm

Triamcinolone Acetonide, USP (TR107)	0.06gm
Ketoconazole, USP (K1149)	0.6gm
Gentamicin, USP	0.18gm
Mineral Oil, Heavy, Oil, USP (MI500)	XmL
Lanolin, USP (LA109)	Q.S. 60gm

SUGGESTED COMPOUNDING PROCEDURES

1. Calculate the required quantity of each ingredient for the amount to be prepared
2. Accurately weigh and/or measure each ingredient
3. Triturate all the dry ingredients together in mortar and pestle
4. Use enough Mineral oil to form a paste and continue to triturate
5. Geometrically mix in Lanolin until homogenous
6. Use of an Ointment mill maybe necessary to reduce particle size
7. Package and label; refrigerate after compounding
 - a. color
 - b. texture
 - c. container
 - d. Label - auxiliary labels, storage, BUD, compounded medication

Store in air-tight container, room temperature

No claims are made as to the safety or efficacy of this preparation. This formulation is provided solely at the unsolicited request of the pharmacist.

Beyond-Use Dates of preparations are conservative estimates from reference books, peer-reviewed literature, and intended duration of therapy, formulation from commercially available products, organoleptic observations and current USP guidelines. Compounders may have stability studies performed by a reputable laboratory if they wish to extend the Beyond-Use Date. It is recommended that you follow USP <795> recommendations for potency testing.

Beyond-Use Date should be assigned based on the current USP <795> Standards

Precautions:

Precautions should be taken to prevent cross-contamination and exposure of ingredients to the compounder and contamination of the preparation by the compounder. Wear appropriate protective equipment. Use safety enclosures (hoods) when weighing and mixing.

11/18 JD