



SUGGESTED FORMULA

Vanillin 2% W/V Oral Flavoring Solution

Version number: 1.0

Volume: 100 mL

Vanillin, NF (VA130)

2gm

Dehydrated Alcohol, USP (ET107)

Q.S. 100mL

***Wear appropriate personal protective equipment**

SUGGESTED COMPOUNDING PROCEDURES

1. Calculate the required quantity of each ingredient for the total amount to be prepared
2. Accurately weigh and/or measure each ingredient
3. Dissolve the Vanillin powder in the alcohol and complete the volume in a graduated cylinder
4. Package in a glass bottle and label. Protect from light
5. Suggested quality assessments:
 - a. Clarity
 - b. Color
 - c. Label (s)- auxiliary labels (external use), storage, BUD

*This is a concentrated Flavoring Solution, dilute prior to use

Store in air-tight container.

Store Refrigerated (2-8°C)

No claims are made as to the safety or efficacy of this preparation. This formulation is provided solely at the unsolicited request of the pharmacist.

Beyond-Use Dates of preparations are conservative estimates by the formulator using reference books, peer-reviewed literature, and intended duration of therapy, formulation from commercially available products, organoleptic observations and current USP guidelines. Compounders may have stability studies performed by a reputable laboratory if they wish to extend the Beyond-Use Date.

It is recommended that you follow USP <795> recommendations for potency testing.

Beyond-Use Date is estimated to be 60 days refrigerated per CompoundingToday.com Formula #1125

Precautions should be taken to prevent cross-contamination and exposure of ingredients to the compounder and contamination of the preparation by the compounder. Wear appropriate protective equipment. Use safety enclosures (hoods) when weighing and mixing.

5/18 JD