

**FIFRA Section 2(ee)
Recommendation**

For aerial application on leafy vegetables and cole crops in CA and AZ only

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U.S. Environmental Protection Agency.

Please contact your Certis USA regional sales manager to determine the specific requirements for FIFRA 2(ee) recommendations in your respective state.

Seduce®

INSECT BAIT

EPA Reg. No. 67702-25-70051

ACTIVE INGREDIENT: Spinosad (a mixture of spinosyn A and spinosyn D):	0.07%
OTHER INGREDIENTS:	99.93%
TOTAL:	100.00%

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

AERIAL APPLICATION

Use appropriate spreading/application equipment to ensure thorough coverage of the treatment area. Apply when wind conditions are favorable and will limit offsite drift. Aerial applications count against the maximum number of yearly applications for spinosad, either by ground or air. Do not apply more than the maximum number of yearly applications specified on the EPA-registered label.

APPLICATION RATE

Apply 20-44 lb/acre.

CROPS

Cole Crops (broccoli, broccoli raab, Brussels sprouts, cabbage, cauliflower, cavalo, Chinese broccoli, Chinese cabbage (bok choy), Chinese cabbage (napa), Chinese mustard cabbage (gai choy), collards, kale, kohlrabi, izuna, mustard greens, mustard spinach, rape greens)

Leafy Vegetables (except Brassica) and leaves of root and tuber and legume vegetables (arugula, beets, celery, peas, lettuce, rhubarb, spinach, swiss chard), turnip greens, and watercress

PESTS

Earwigs, cutworms

MAXIMUM APPLICATIONS

Do not make more than 6 applications per calendar year.

APPLICATION INTERVAL

Do not make applications less than 4 days apart.

PRE-HARVEST INTERVAL

Do not apply within one day of harvest.

OTHER RESTRICTIONS

For cole crops: Do not apply more than a total of 0.45 lb ai spinosad (including foliar uses) per acre per crop. Do not apply to seedling cole crops grown for transplant.

For leafy vegetables: Do not apply more than a total of 0.45 lb ai spinosad (including foliar uses) per acre per crop. Do not apply to seedling leafy crops grown for transplant within a greenhouse or shade house.

All applicable directions, restrictions, and precautions on the EPA-registered labels are to be followed.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and observe the cautionary statements plus all other information appearing on the product labels. This recommendation must be in the possession of the user at the time of pesticide application.