

HIGHLIGHTS OF PRESCRIBING INFORMATION
These highlights do not include all the information needed to use TERIPARATIDE INJECTION safely and effectively. See full prescribing information for TERIPARATIDE INJECTION.

TERIPARATIDE injection, for subcutaneous use
Initial U.S. Approval: 1987

WARNING: POTENTIAL RISK OF OSTEOSARCOMA
See full prescribing information for complete boxed warning.

- In rats, teriparatide caused an increase in the incidence of osteosarcoma, a malignant bone tumor. (5.1, 13.1)
- Because of the uncertain relevance of the rat osteosarcoma finding to humans, prescribe teriparatide injection only for patients for whom potential benefits outweigh potential risk. (5.1)
- Teriparatide injection should not be prescribed for patients at increased baseline risk for osteosarcoma (e.g., those with Paget's disease of bone or unexplained elevations of alkaline phosphatase, pediatric and young adult patients with open epiphyses, or prior external beam or implant radiation therapy involving the skeleton). (5.1)

INDICATIONS AND USAGE
Teriparatide injection is a parathyroid hormone analog (PTH 1-34) indicated for:

- Treatment of postmenopausal women with osteoporosis at high risk for fracture (1.1)
- Increase of bone mass in men with primary or hypogonadal osteoporosis at high risk for fracture (1.2)
- Treatment of men and women with osteoporosis associated with sustained systemic glucocorticoid therapy at high risk for fracture (1.3)

DOSAGE AND ADMINISTRATION
Recommended dose is 20 mcg subcutaneously once a day (2.1, 2.2, 2.3)

- Administer as a subcutaneous injection into the thigh or abdominal wall (2.4)
- Administer initially under circumstances in which the patient can sit or lie down if symptoms of orthostatic hypotension occur (2.4)
- Use of the drug for more than 2 years during a patient's lifetime is not recommended (2.5)

DOSAGE FORMS AND STRENGTHS
Injection: 620 mcg/2.48 mL (250 mcg/mL) in single-patient-use pen containing 28 daily doses of 20 mcg (3)

CONTRAINDICATIONS

- Patients with hypersensitivity to teriparatide or to any of its excipients (4)

WARNINGS AND PRECAUTIONS

- Patients with Paget's disease of bone, pediatric and young adult patients with open epiphyses, and patients with prior external beam or implant radiation involving the skeleton: Should not be treated with teriparatide injection (5.1, 8.4)

Revised: 11/2019

FULL PRESCRIBING INFORMATION: CONTENTS*
WARNING: POTENTIAL RISK OF OSTEOSARCOMA
1 INDICATIONS AND USAGE
1.1 Treatment of Postmenopausal Women with Osteoporosis at High Risk for Fracture
1.2 Increase of Bone Mass in Men with Primary or Hypogonadal Osteoporosis at High Risk for Fracture
1.3 Treatment of Men and Women with Glucocorticoid-Induced Osteoporosis at High Risk for Fracture
2 DOSAGE AND ADMINISTRATION
2.1 Treatment of Postmenopausal Women with Osteoporosis at High Risk for Fracture
2.2 Increase of Bone Mass in Men with Primary or Hypogonadal Osteoporosis at High Risk for Fracture
2.3 Treatment of Men and Women with Glucocorticoid-Induced Osteoporosis at High Risk for Fracture
2.4 Administration
2.5 Treatment Duration
3 DOSAGE FORMS AND STRENGTHS
4 CONTRAINDICATIONS
5 WARNINGS AND PRECAUTIONS
5.1 Osteosarcoma
5.2 Treatment Duration
5.3 Bone Metastases and Skeletal Malignancies
5.4 Metabolic Bone Diseases
5.5 Hypercalcemia and Hypercalcemic Disorders
5.6 Urolithiasis or Pre-existing Hypercalciuria
5.7 Orthostatic Hypotension
5.8 Drug Interactions
6 ADVERSE REACTIONS
6.1 Clinical Trials Experience
6.2 Postmarketing Experience
6.3 Immunogenicity
7 DRUG INTERACTIONS
7.1 Digoxin
7.2 Hydrochlorothiazide
7.3 Furosemide

USE IN SPECIFIC POPULATIONS
WARNING: POTENTIAL RISK OF OSTEOSARCOMA
In male and female rats, teriparatide caused an increase in the incidence of osteosarcoma (a malignant bone tumor) that was dependent on dose and treatment duration. The effect was observed at systemic exposures to teriparatide ranging from 3 to 60 times the exposure in humans given a 20 mcg dose. Because of the uncertain relevance of the rat osteosarcoma finding to humans, prescribe teriparatide injection only for patients for whom the potential benefits are considered to outweigh the potential risk. Teriparatide injection should not be prescribed for patients who are at increased baseline risk for osteosarcoma (including those with Paget's disease of bone or unexplained elevations of alkaline phosphatase, pediatric and young adult patients with open epiphyses, or prior external beam or implant radiation therapy involving the skeleton) [see Warnings and Precautions (5.1), Adverse Reactions (6.2), and Nonclinical Toxicology (13.1)].

1 INDICATIONS AND USAGE
1.1 Treatment of Postmenopausal Women with Osteoporosis at High Risk for Fracture
Teriparatide injection is indicated for the treatment of postmenopausal women with osteoporosis at high risk for fracture, defined as a history of osteoporotic fracture, multiple risk factors for fracture, or patients who have failed or are intolerant to other available osteoporosis therapy. In postmenopausal women with osteoporosis, teriparatide injection reduces the risk of vertebral and nonvertebral fractures [see Clinical Studies (14.1)].
1.2 Increase of Bone Mass in Men with Primary or Hypogonadal Osteoporosis at High Risk for Fracture
Teriparatide injection is indicated to increase bone mass in men with primary or hypogonadal osteoporosis at high risk for fracture, defined as a history of osteoporotic fracture, multiple risk factors for fracture, or patients who have failed or are intolerant to other available osteoporosis therapy [see Clinical Studies (14.2)].
1.3 Treatment of Men and Women with Glucocorticoid-Induced Osteoporosis at High Risk for Fracture
Teriparatide injection is indicated for the treatment of men and women with osteoporosis associated with sustained systemic glucocorticoid therapy (daily dosage equivalent to 5 mg or greater of prednisone) at high risk for fracture, defined as a history of osteoporotic fracture, multiple risk factors for fracture, or patients who have failed or are intolerant to other available osteoporosis therapy [see Clinical Studies (14.3)].

2 DOSAGE AND ADMINISTRATION
2.1 Treatment of Postmenopausal Women with Osteoporosis at High Risk for Fracture
The recommended dose is 20 mcg subcutaneously once a day.
2.2 Increase of Bone Mass in Men with Primary or Hypogonadal Osteoporosis at High Risk for Fracture
The recommended dose is 20 mcg subcutaneously once a day.
2.3 Treatment of Men and Women with Glucocorticoid-Induced Osteoporosis at High Risk for Fracture
The recommended dose is 20 mcg subcutaneously once a day.
2.4 Administration
* Teriparatide injection should be administered as a subcutaneous injection into the thigh or abdominal wall. There are no data available on the safety or efficacy of intravenous or intramuscular injection of teriparatide.
* Teriparatide injection should be administered initially under circumstances in which the patient can sit or lie down if symptoms of orthostatic hypotension occur [see Warnings and Precautions (5.7)].
* Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit. Teriparatide injection is a clear and colorless liquid. Do not use if solid particles appear or if the solution is cloudy or colored.
* Patients and caregivers who administer teriparatide injection should receive appropriate training and instruction on the proper use of the teriparatide injection delivery device from a qualified health professional [see Patient Counseling Information (17.5)].

2.5 Treatment Duration
The safety and efficacy of teriparatide have not been evaluated beyond 2 years of treatment. Consequently, use of the drug for more than 2 years during a patient's lifetime is not recommended.
3 DOSAGE FORMS AND STRENGTHS
Injection: 620 mcg/2.48 mL (250 mcg/mL)
Colorless solution in a single-patient-use pen containing 28 daily doses of 20 mcg.
4 CONTRAINDICATIONS
Teriparatide injection is contraindicated in patients with:

- Hypersensitivity to teriparatide or to any of its excipients. Reactions have included angioedema and anaphylaxis [see Adverse Reactions (6.2)].

5 WARNINGS AND PRECAUTIONS
5.1 Osteosarcoma
In male and female rats, teriparatide caused an increase in the incidence of osteosarcoma (a malignant bone tumor) that was dependent on dose and treatment duration [see Boxed Warning and Nonclinical Toxicology (13.1)]. Teriparatide injection should not be prescribed for patients at increased baseline risk of osteosarcoma.
These include:

- Paget's disease of bone. Unexplained elevations of alkaline phosphatase may indicate Paget's disease of bone.
- Pediatric and young adult patients with open epiphyses.
- Prior external beam or implant radiation therapy involving the skeleton.

5.2 Treatment Duration
The safety and efficacy of teriparatide have not been evaluated beyond 2 years of treatment. Consequently, use of the drug for more than 2 years during a patients' lifetime is not recommended.
5.3 Bone Metastases and Skeletal Malignancies
Patients with bone metastases or a history of skeletal malignancies should not be treated with teriparatide injection.
5.4 Metabolic Bone Diseases
Patients with metabolic bone diseases other than osteoporosis should not be treated with teriparatide injection.
5.5 Hypercalcemia and Hypercalcemic Disorders
Teriparatide has not been studied in patients with pre-existing hypercalcemia. These patients should not be treated with teriparatide because of the possibility of exacerbating hypercalcemia. Patients known to have an underlying hypercalcemic disorder, such as primary hyperparathyroidism, should not be treated with teriparatide injection.
5.6 Urolithiasis or Pre-existing Hypercalciuria
In clinical trials, the frequency of urolithiasis was similar in patients treated with teriparatide and placebo. However, teriparatide has not been studied in patients with active urolithiasis. If active urolithiasis or pre-existing hypercalciuria are suspected, measurement of urinary calcium excretion should be considered. Teriparatide injection should be used with caution in patients with active or recent urolithiasis because of the potential to exacerbate this condition.

5.7 Orthostatic Hypotension
Teriparatide injection should be administered initially under circumstances in which the patient can sit or lie down if symptoms of orthostatic hypotension occur. In short-term clinical pharmacology studies with teriparatide, transient episodes of symptomatic orthostatic hypotension were observed in 5% of patients. Typically, an event began within 4 hours of dosing and spontaneously resolved within a few minutes to a few hours. When transient orthostatic hypotension occurred, it happened within the first several doses, it was relieved by placing the person in a reclining position, and it did not preclude continued treatment.
5.8 Drug Interactions
Hypercalcemia may predispose patients to digitalis toxicity. Because teriparatide transiently increases serum calcium, patients receiving digoxin should use teriparatide injection with caution [see Drug Interactions (7.1) and Clinical Pharmacology (12.2)].

6 ADVERSE REACTIONS
6.1 Clinical Trials Experience
Because clinical studies are conducted under widely varying conditions, adverse reaction rates observed in the clinical studies of a drug cannot be directly compared to rates in the clinical studies of another drug and may not reflect the rates observed in practice.
Treatment of Osteoporosis in Men and Postmenopausal Women
The safety of teriparatide in the treatment of osteoporosis in men and postmenopausal women was assessed in two randomized, double-blind, placebo-controlled trials of 1382 patients (21% men, 79% women) aged 28 to 86 years (mean 67 years). The median durations of the trials were 11 months for men and 19 months for women, with 691 patients exposed to teriparatide and 691 patients to placebo. All patients received 1000 mg of calcium plus at least 400 IU of vitamin D supplementation per day.
The incidence of all cause mortality was 1% in the teriparatide group and 1% in the placebo group. The incidence of serious adverse events was 16% in teriparatide patients and 19% in placebo patients. Early discontinuation due to adverse events occurred in 7% of teriparatide patients and 6% of placebo patients.

Table 1 lists adverse events from the two principal osteoporosis trials in men and postmenopausal women that occurred in ≥2% of teriparatide-treated and more frequently than placebo-treated patients.

Table 1. Percentage of Patients with Adverse Events Reported by at Least 2% of Teriparatide-Treated Patients and in More Teriparatide-Treated Patients than Placebo-Treated Patients from the Two Principal Osteoporosis Trials in Women and Men (Adverse Events are Shown Without Attribution of Causality)

	Teriparatide N=691 (%)	Placebo N=691 (%)
Event Classification		
Body as a Whole		
Pain	21.3	20.5
Headache	7.5	7.4
Asthenia	8.7	6.8
Neck pain	3.0	2.7
Cardiovascular		
Hypertension	7.1	6.8
Angina pectoris	2.5	1.6
Syncope	2.6	1.4
Digestive System		
Nausea	8.5	6.7
Constipation	5.4	4.5
Diarrhea	5.1	4.6
Dyspepsia	5.2	4.1
Vomiting	3.0	2.3
Gastrointestinal disorder	2.3	2.0
Tooth disorder	2.0	1.3
Musculoskeletal		
Arthralgia	10.1	8.4
Leg cramps	2.6	1.3
Nervous System		
Dizziness	8.0	5.4
Depression	4.1	2.7
Insomnia	4.3	3.6
Vertigo	3.8	2.7
Respiratory System		
Rhinitis	9.6	8.8
Cough increased	6.4	5.5
Pharyngitis	5.5	4.8
Dyspnea	3.6	2.6
Pneumonia	3.9	3.3
Skin and Appendages		
Rash	4.9	4.5
Sweating	2.2	1.7

Laboratory Findings
Serum Calcium — Teriparatide transiently increased serum calcium, with the maximal effect observed at approximately 4 to 6 hours post-dose. Serum calcium measured at least 16 hours post-dose was not different from pretreatment levels. In clinical trials, the frequency of at least 1 episode of transient hypercalcemia in the 4 to 6 hours after teriparatide administration was increased from 2% of women and none of the men treated with placebo to 11% of women and 6% of men treated with teriparatide. The number of patients treated with teriparatide whose transient hypercalcemia was verified on consecutive measurements was 3% of women and 1% of men.
Urinary Calcium — Teriparatide increased urinary calcium excretion, but the frequency of hypercalciuria in clinical trials was similar for patients treated with teriparatide and placebo [see Clinical Pharmacology (12.2)].
Serum Uric Acid — Teriparatide increased serum uric acid concentrations. In clinical trials, 3% of teriparatide patients had serum uric acid concentrations above the upper limit of normal compared with 1% of placebo patients. However, the hyperuricemia did not result in an increase in gout, arthralgia, or urolithiasis.

Renal Function — No clinically important adverse renal effects were observed in clinical studies. Assessments included creatinine clearance; measurements of blood urea nitrogen (BUN), creatinine, and electrolytes in serum; urine specific gravity and pH; and examination of urine sediment.

Studies in Men and Women with Glucocorticoid-Induced Osteoporosis
The safety of teriparatide in the treatment of men and women with glucocorticoid-induced osteoporosis was assessed in a randomized, double-blind, active-controlled trial of 428 patients (19% men, 81% women) aged 22 to 89 years (mean 57 years) treated with ≥5 mg per day prednisone or equivalent for a minimum of 3 months. The duration of the trial was 18 months with 214 patients exposed to teriparatide and 214 patients exposed to oral daily bisphosphonate (active control). All patients received 1000 mg of calcium plus 800 IU of vitamin D supplementation per day.
The incidence of all cause mortality was 4% in the teriparatide group and 6% in the active control group. The incidence of serious adverse events was 21% in teriparatide patients and 18% in active control patients, and included pneumonia (3% teriparatide, 1% active control). Early discontinuation because of adverse events occurred in 15% of teriparatide patients and 12% of active control patients, and included dizziness (2% teriparatide, 0% active control).
Adverse events reported at a higher incidence in the teriparatide group and with at least a 2% difference in teriparatide-treated patients compared with active control-treated patients were: nausea (14%, 7%), gastritis (7%, 3%), pneumonia (6%, 3%), dyspnea (6%, 3%), insomnia (5%, 1%), anxiety (4%, 1%), and herpes zoster (3%, 1%), respectively.

6.2 Postmarketing Experience
The following adverse reactions have been identified during post-approval use of teriparatide. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

- Osteosarcoma: Cases of bone tumor and osteosarcoma have been reported rarely in the postmarketing period. The causality to teriparatide use is unclear. Long term osteosarcoma surveillance studies are ongoing [see Warnings and Precautions (5.1)].
- Hypercalcemia: Hypercalcemia greater than 13.0 mg/dL has been reported with teriparatide use.
- Adverse events reported since market introduction that were temporally (but not necessarily causally) related to teriparatide therapy include the following:
 - Allergic Reactions: Anaphylactic reactions, drug hypersensitivity, angioedema, urticaria
 - Investigations: Hyperuricemia
 - Respiratory System: Acute dyspnea, chest pain
 - Musculoskeletal: Muscle spasms of the leg or back
 - Other: Injection site reactions including injection site pain, swelling and bruising; oro-facial edema

6.3 Immunogenicity
As with all peptides, there is potential for immunogenicity. The detection of antibody formation is highly dependent on the sensitivity and specificity of the assay. Additionally, the observed incidence of antibody (including neutralizing antibody) positivity in an assay may be influenced by several factors, including assay methodology, sample handling, timing of sample collection, concomitant medications, and underlying disease. For these reasons, comparison of the incidence of antibodies to teriparatide in the study described below with the incidence of antibodies in other studies or to other teriparatide products may be misleading.
The immunogenicity profile of teriparatide injection was evaluated in a 24-week randomized trial comparing the effects of teriparatide injection 20 mcg daily with another teriparatide product 20 mcg daily. In this trial, 2.2% (2/90) subjects who received teriparatide injection and 2.2% (2/91) subjects who received the other teriparatide product had detectable antibodies to teriparatide and one out of the two teriparatide injection treated patients developed neutralizing antibodies to teriparatide.

7 DRUG INTERACTIONS
7.1 Digoxin
A single teriparatide dose did not alter the effect of digoxin on the systolic time interval (from electrocardiographic Q-wave onset to aortic valve closure, a measure of digoxin's calcium-mediated cardiac effect). However, because teriparatide may transiently increase serum calcium, teriparatide injection should be used with caution in patients taking digoxin [see Warnings and Precaution (5.8) and Clinical Pharmacology (12.2)].
7.2 Hydrochlorothiazide
The coadministration of hydrochlorothiazide 25 mg with teriparatide did not affect the serum calcium response to teriparatide 40 mcg. The effect of coadministration of a higher dose of hydrochlorothiazide with teriparatide on serum calcium levels has not been studied [see Clinical Pharmacology (12.2)].
7.3 Furosemide
Coadministration of intravenous furosemide (20 to 100 mg) with teriparatide 40 mcg in healthy people and patients with mild, moderate, or severe renal impairment (CrCl 13 to 72 mL/min) resulted in small increases in the serum calcium (2%) and 24-hour urine calcium (37%) responses to teriparatide that did not appear to be clinically important [see Clinical Pharmacology (12.2)].

8 USE IN SPECIFIC POPULATIONS
8.1 Pregnancy
Risk Summary
There are no available data on teriparatide injection use in pregnant women to evaluate for drug-associated risk of major birth defects, miscarriage, or adverse maternal or fetal outcomes. Consider discontinuing teriparatide injection when pregnancy is recognized.
In animal reproduction studies, teriparatide increased skeletal deviations and variations in mouse offspring at subcutaneous doses equivalent to more than 60 times the recommended 20 mcg human daily dose (based on body surface area, mcg/m²), and produced mild growth retardation and reduced motor activity in rat offspring at subcutaneous doses equivalent to more than 120 times the human dose (see Data).
The background risk of major birth defects and miscarriage for the indicated population is unknown. The background risk in the US general population of major birth defects is 2% to 4% and of miscarriage is 15% to 20% of clinically recognized pregnancies.
Data
Animal Data
In animal reproduction studies, pregnant mice received teriparatide during organogenesis at subcutaneous doses 8 to 267 times the human dose (based on body surface area, mcg/m²). At doses ≥60 times the human dose, the fetuses showed an increased incidence of skeletal deviations or variations (interrupted rib, extra vertebra or rib). When pregnant rats received subcutaneous teriparatide during organogenesis at subcutaneous doses 16 to 540 times the human dose, the fetuses showed no abnormal findings.
In a perinatal/postnatal study in pregnant rats dosed subcutaneously from organogenesis through lactation, mild growth retardation was observed in female offspring at doses ≥120 times the human dose. Mild growth retardation in male offspring and reduced motor activity in both male and female offspring were observed at maternal doses of 540 times the human dose. There were no developmental or reproductive effects in mice or rats at doses 8 or 16 times the human dose, respectively.

8.2 Lactation
Risk Summary
It is not known whether teriparatide is excreted in human milk, affects human milk production, or has effects on the breast fed infant. Because of the potential for osteosarcoma shown with teriparatide in animal studies, advise patients that breastfeeding is not recommended during treatment with teriparatide injection [see Warnings and Precautions (5.1)].
8.4 Pediatric Use
The safety and efficacy of teriparatide have not been established in any pediatric population. Teriparatide injection should not be prescribed in patients at an increased baseline risk of osteosarcoma which include pediatric and young adult patients with open epiphyses. Therefore, teriparatide injection is not indicated for use in pediatric or young adult patients with open epiphyses [see Warnings and Precautions (5.1)].
8.5 Geriatric Use
Of the patients receiving teriparatide in the osteoporosis trial of 1637 postmenopausal women, 75% were 65 years of age and over and 23% were 75 years of age and over. Of the patients receiving teriparatide in the osteoporosis trial of 437 men, 39% were 65 years of age and over and 13% were 75 years of age and over. No overall differences in safety or effectiveness were observed between these subjects and younger subjects, and other reported clinical experience has not identified differences in responses between the elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out.

8.6 Hepatic Impairment
No studies have been performed in patients with hepatic impairment [see Clinical Pharmacology (12.3)].
8.7 Renal Impairment
In 5 patients with severe renal impairment (CrCl <30 mL/min), the AUC and T_{1/2} of teriparatide were increased by 73% and 77%, respectively. Maximum serum concentration of teriparatide was not increased [see Clinical Pharmacology (12.3)].
10 OVERDOSAGE
Incidents of overdose in humans have not been reported in clinical trials. Teriparatide has been administered in single doses of up to 100 mcg and in repeated doses of up to 60 mcg/day for 6 weeks. The effects of overdose that might be expected include a delayed hypercalcemic effect and risk of orthostatic hypotension. Nausea, vomiting, dizziness, and headache might also occur.

Medication Guide

Teriparatide (ter i par a tide) Injection

Read this Medication Guide before you start taking teriparatide injection and each time you get a refill. There may be new information. Also, read the *User Manual* that comes with the teriparatide injection device (pen) for information on how to use the device to inject your medicine the right way. This Medication Guide does not take the place of talking with your healthcare provider about your medical condition or your treatment.

What is the most important information I should know about teriparatide injection?

- Possible bone cancer.** During drug testing, the medicine in teriparatide injection caused some rats to develop a bone cancer called osteosarcoma. In people, osteosarcoma is a serious but rare cancer. Osteosarcoma has rarely been reported in people who have taken teriparatide injection. It is not known if people who take teriparatide injection have a higher chance of getting osteosarcoma.
- You should not take teriparatide injection for more than 2 years over your lifetime.

What is teriparatide injection?

- Teriparatide injection is a prescription medicine that is like a hormone made by the body called parathyroid hormone or PTH.
- Teriparatide injection may help to form new bone, increase bone mineral density and bone strength.
- Teriparatide injection can lessen the number of fractures of the spine and other bones in postmenopausal women with osteoporosis.
- The effect on fractures has not been studied in men.

- Teriparatide injection is used in both men and postmenopausal women with osteoporosis who are at high risk for having fractures. Teriparatide injection can be used by people who have had a fracture related to osteoporosis, or who have several risk factors for fracture, or who can not use other osteoporosis treatments.
- Teriparatide injection is used in both men and women with osteoporosis due to use of glucocorticoid medicines, such as prednisone, for several months, who are at high risk for having broken bones (fractures). These include men and women with either a history of broken bones, who have several risk factors for fracture, or who can not use other osteoporosis treatments.

It is not known if teriparatide injection is safe and effective in children.

Teriparatide injection should not be used in children and young adults whose bones are still growing.

Who should not use teriparatide injection?

Do not use teriparatide injection if you are allergic to any of the ingredients in teriparatide injection. See the end of this Medication Guide for a complete list of the ingredients in teriparatide injection.

Before you take teriparatide injection, tell your healthcare provider if you:

- have the condition listed in the section "Who should not use teriparatide injection?"
- have Paget's disease or other bone disease
- have cancer in your bones

- have trouble injecting yourself and do not have someone who can help you

- are a child or young adult whose bones are still growing
- have or have had kidney stones
- have had radiation therapy

- have or had too much calcium in your blood
- have any other medical conditions

- are pregnant or thinking about becoming pregnant. It is not known if teriparatide injection will harm your unborn baby.

- are breast-feeding or plan to breast-feed. You should not breast-feed while taking teriparatide injection.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Your healthcare provider needs this information to help keep you from taking teriparatide injection with other medicines that may harm you.

Especially tell your doctor if you take medicines that contain digoxin (Digoxin*, Lanoxicaps*, Lanoxin*).

How should I use teriparatide injection?

- Inject teriparatide injection one time each day in your thigh or abdomen (lower stomach area). Talk to a healthcare provider about how to rotate injection sites.

- Before you try to inject teriparatide injection yourself, a healthcare provider should teach you how to use the teriparatide injection delivery device to give your injection the right way.
- Read the detailed *User Manual*.

- You can take teriparatide injection with or without food or drink.

- The teriparatide injection delivery device has enough medicine for 28 days. It is set to give a 20 microgram dose of medicine each day. Do not inject all the medicine in the teriparatide injection delivery device at any one time.

- Do not transfer the medicine from the teriparatide injection delivery device to a syringe. This can result in taking the wrong dose of teriparatide injection. If you do not have pen needles to use with your teriparatide injection delivery device, talk with your healthcare provider.

- Teriparatide injection should look clear and colorless. Do not use teriparatide injection if it has particles in it, or if it is cloudy or colored.

- Inject teriparatide injection right away after you take the delivery device out of the refrigerator.

- After each use, safely remove the needle, recap the delivery device, and put it back in the refrigerator right away.

- You can take teriparatide injection at any time of the day. To help you remember to take teriparatide injection, take it at about the same time each day.

- If you forget or cannot take teriparatide injection at your usual time, take it as soon as you can on that day. Do not take more than one injection in the same day.

- If you take more teriparatide injection than prescribed, call your healthcare provider. If you take too much teriparatide injection, you may have nausea, vomiting, weakness, or dizziness.

- Follow your healthcare provider's instructions about other ways you can help your osteoporosis, such as exercise, diet, and reducing or stopping your use of tobacco and alcohol. If your healthcare provider recommends calcium and vitamin D supplements, you can take them at the same time you take teriparatide injection.

What are the possible side effects of teriparatide injection?

Teriparatide injection can cause serious side effects, including:

- See **“What is the most important information I should know about teriparatide injection?”**
- Decrease in blood pressure when you change positions.** Some people feel dizzy, get a fast heartbeat, or feel faint right after the first few doses. This usually happens within 4 hours of taking teriparatide injection and goes away within a few hours. For the first few doses, take your injections of teriparatide injection in a place where you can sit or lie down right away if you get these symptoms. If your symptoms get worse or do not go away, stop taking teriparatide injection and call your healthcare provider.
- Increased calcium in your blood.** Tell your healthcare provider if you have nausea, vomiting, constipation, low energy, or muscle weakness. These may be signs that there is too much calcium in your blood.

Common side effects of teriparatide injection include:

- nausea
- joint aches
- pain

Your healthcare provider may take samples of blood and urine during treatment to check your response to teriparatide injection. Also, your healthcare provider may ask you to have follow-up tests of bone mineral density.

Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all the possible side effects of teriparatide injection. For more information, ask your doctor or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store teriparatide injection?

- Keep your teriparatide injection delivery device in the refrigerator between 36°F to 46°F (2°C to 8°C).
- Do not freeze the teriparatide injection delivery device. Do not use teriparatide injection if it has been frozen.
- Do not use teriparatide injection after the expiration date printed on the delivery device and packaging.
- Throw away the teriparatide injection delivery device after 28 days even if it has medicine in it (see the User Manual).

Keep teriparatide injection and all medicines out of the reach of children.

General information about teriparatide injection

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use teriparatide injection for a condition for which it was not prescribed. Do not give teriparatide injection to other people, even if they have the same condition you have.

This Medication Guide summarizes the most important information about teriparatide injection. If you would like more information, talk with your healthcare provider. You can ask your pharmacist or healthcare provider for information about teriparatide injection that is written for healthcare professionals. For more information, call Alvogin, Inc. at 1-866-770-3024.

What are the ingredients in teriparatide injection?

Active ingredient: teriparatide

Inactive ingredients: glacial acetic acid, sodium acetate, mannitol, metacresol, and water for injection.

What is Osteoporosis?

Osteoporosis is a disease in which the bones become thin and weak, increasing the chance of having a broken bone. Osteoporosis usually causes no symptoms until a fracture happens. The most common fractures are in the spine (backbone). They can shorten height, even without causing pain. Over time, the spine can become curved or deformed and the body bent over. Fractures from osteoporosis can also happen in almost any bone in the body, for example, the wrist, rib, or hip. Once you have had a fracture, the chance for more fractures greatly increases.

The following risk factors increase your chance of getting fractures from osteoporosis:

- past broken bones from osteoporosis
- very low bone mineral density (BMD)
- frequent falls
- limited movement, such as using a wheelchair
- medical conditions likely to cause bone loss, such as some kinds of arthritis
- taking steroid medicines called glucocorticoids, such as prednisone
- other medicines that may cause bone loss, for example: seizure medicines (such as phenytoin), blood thinners (such as heparin), and high doses of vitamin A

*The brands listed are trademarks of their respective owners and are not trademarks of Alvogin, Inc.

Distributed by:

Alvogin, Inc.
Morristown, NJ 07960 USA

©2017 Alvogin, Inc. All rights reserved.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Revised: November 2019


In postmarketing spontaneous reports, there have been cases of medication errors in which the entire contents (up to 800 mcg) of the teriparatide delivery device (pen) have been administered as a single dose. Transient events reported have included nausea, weakness/lethargy and hypotension. In some cases, no adverse events occurred as a result of the overdose. No fatalities associated with overdose have been reported.

Overdose Management — There is no specific antidote for teriparatide. Treatment of suspected overdose should include discontinuation of teriparatide injection, monitoring of serum calcium and phosphorus, and implementation of appropriate supportive measures, such as hydration.

1. DESCRIPTION

Teriparatide injection is a recombinant human parathyroid hormone analog (PTH 1-34). It has an identical sequence to the 34 N-terminal amino acids (the biologically active region) of the 84-amino acid human parathyroid hormone.

The molecular formula of teriparatide is C₁₂₁H₂₂₈N₃₂O₅₂S₂ and the molecular weight is 4117.8 daltons. Its amino acid sequence is shown below:



Teriparatide is manufactured using a strain of *Pseudomonas fluorescens* modified by recombinant DNA technology. Teriparatide injection is supplied as a sterile, colorless, clear, isotonic solution in a glass cartridge which is pre-assembled into a single-patient-use delivery device (pen) for subcutaneous injection. Each pen is filled to deliver 2.48 mL. Each mL contains 250 mcg teriparatide (as a free base), 0.41 mg glacial acetic acid, 0.10 mg sodium acetate, 45.4 mg mannitol, 3 mg metacresol, and water for injection. The drug product is a pH 4.0 solution.

Each cartridge, pre-assembled into a delivery device, delivers 20 mcg of teriparatide per dose each day for up to 28 days.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Endogenous 84-amino acid parathyroid hormone (PTH) is the primary regulator of calcium and phosphate metabolism in bone and kidney. Physiological actions of PTH include regulation of bone metabolism, renal tubular reabsorption of calcium and phosphate, and intestinal calcium absorption. The biological actions of PTH and teriparatide are mediated through binding to specific high-affinity cell-surface receptors. Teriparatide and the 34 N-terminal amino acids of PTH bind to these receptors with the same affinity and have the same physiological actions on bone and kidney. Teriparatide is not expected to accumulate in bone or other tissues.

The skeletal effects of teriparatide depend upon the pattern of systemic exposure. Once-daily administration of teriparatide stimulates new bone formation on trabecular and cortical (periosteal) bone surfaces by preferential stimulation of osteoblastic activity over osteoclastic activity. In monkey studies, teriparatide improved trabecular microarchitecture and increased bone mass, and strength by stimulating new bone formation in both cancellous and cortical bone. In humans, the anabolic effects of teriparatide manifest as an increase in skeletal mass, an increase in markers of bone formation and resorption, and an increase in bone strength. By contrast, continuous excess of endogenous PTH, as occurs in hyperparathyroidism, may be detrimental to the skeleton because bone resorption may be stimulated more than bone formation.

12.2 Pharmacodynamics

Pharmacodynamics in Men and Postmenopausal Women with Osteoporosis

Effects on Mineral Metabolism — Teriparatide affects calcium and phosphorus metabolism in a pattern consistent with the known actions of endogenous PTH (e.g., increases serum calcium and decreases serum phosphorus).

Serum Calcium Concentrations — When teriparatide 20 mcg is administered once daily, the serum calcium concentration increases transiently, beginning approximately 2 hours after dosing and reaching a maximum concentration between 4 and 6 hours (median increase, 0.4 mg/dL). The serum calcium concentration begins to decline approximately 6 hours after dosing and returns to baseline by 16 to 24 hours after each dose.

In a clinical study of postmenopausal women with osteoporosis, the median peak serum calcium concentration measured 4 to 6 hours after dosing with teriparatide 20 mcg was 2.42 mmol/L (9.63 mg/dL) at 12 months. The peak serum calcium remained below 2.76 mmol/L (11.0 mg/dL) in >99% of women at each visit. Sustained hypercalcemia was not observed.

In this study, 11.1% of women treated with teriparatide had at least 1 serum calcium value above the upper limit of normal [2.64 mmol/L (10.6 mg/dL)] compared with 1.5% of women treated with placebo. The percentage of women treated with teriparatide whose serum calcium was above the upper limit of normal on consecutive 4- to 6-hour post-dose measurements was 3.0% compared with 0.2% of women treated with placebo. In these women, calcium supplements and/or teriparatide doses were reduced. The timing of these dose reductions was at the discretion of the investigator. Teriparatide dose adjustments were made at varying intervals after the first observation of increased serum calcium (median 21 weeks). During these intervals, there was no evidence of progressive increases in serum calcium.

In a clinical study of men with either primary or hypogonadal osteoporosis, the effects on serum calcium were similar to those observed in postmenopausal women. The median peak serum calcium concentration measured 4 to 6 hours after dosing with teriparatide was 2.35 mmol/L (9.44 mg/dL) at 12 months. The peak serum calcium remained below 2.76 mmol/L (11.0 mg/dL) in 98% of men at each visit. Sustained hypercalcemia was not observed.

In this study, 6.0% of men treated with teriparatide daily had at least 1 serum calcium value above the upper limit of normal [2.64 mmol/L (10.6 mg/dL)] compared with none of the men treated with placebo. The percentage of men treated with teriparatide whose serum calcium was above the upper limit of normal on consecutive measurements was 1.3% (2 men) compared with none of the men treated with placebo. Although calcium supplements and/or teriparatide doses could have been reduced in these men, only calcium supplementation was reduced [see *Warnings and Precautions (5.5) and Adverse Reactions (6.1)*].

In a clinical study of women previously treated for 18 to 39 months with raloxifene (n=26) or alendronate (n=33), mean serum calcium >12 hours after teriparatide injection was increased by 0.09 to 0.14 mmol/L (0.36 to 0.56 mg/dL), after 1 to 6 months of teriparatide treatment compared with baseline. Of the women pretreated with raloxifene, 3 (11.5%) had a serum calcium >2.76 mmol/L (11.0 mg/dL), and 1 of those pretreated with alendronate, 3 (9.1%) had a serum calcium >2.76 mmol/L (11.0 mg/dL). The highest serum calcium reported was 3.12 mmol/L (12.5 mg/dL). None of the women had symptoms of hypercalcemia. There were no placebo controls in this study.

In the study of patients with glucocorticoid-induced osteoporosis, the effects of teriparatide on serum calcium were similar to those observed in postmenopausal women with osteoporosis not taking glucocorticoids.

Urinary Calcium Excretion — In a clinical study of postmenopausal women with osteoporosis who received 1000 mg of supplemental calcium and at least 400 IU of vitamin D, daily teriparatide increased urinary calcium excretion. The median urinary excretion of calcium was 4.8 mmol/day (190 mg/day) at 6 months and 4.2 mmol/day (170 mg/day) at 12 months. These levels were 0.76 mmol/day (30 mg/day) and 0.3 mmol/day (12 mg/day) higher, respectively, than in women treated with placebo. The incidence of hypercalciuria (>7.5 mmol Ca/day or 300 mg/day) was similar in the women treated with teriparatide or placebo.

In a clinical study of men with either primary or hypogonadal osteoporosis who received 1000 mg of supplemental calcium and at least 400 IU of vitamin D, daily teriparatide had inconsistent effects on urinary calcium excretion. The median urinary excretion of calcium was 5.6 mmol/day (220 mg/day) at 1 month and 5.3 mmol/day (210 mg/day) at 6 months. These levels were 0.5 mmol/day (20 mg/day) higher and 0.2 mmol/day (8 mg/day) lower, respectively, than in men treated with placebo. The incidence of hypercalciuria (>7.5 mmol Ca/day or 300 mg/day) was similar in the men treated with teriparatide or placebo.

Phosphorus and Vitamin D — In single-dose studies, teriparatide produced transient phosphaturia and mild transient reductions in serum phosphorus concentration. However, hypophosphatemia (<0.74 mmol/L or 2.4 mg/dL) was not observed in clinical trials with teriparatide.

In clinical trials of daily teriparatide, the median serum concentration of 1,25-dihydroxyvitamin D was increased at 12 months by 19% in women and 14% in men, compared with baseline. In the placebo group, this concentration decreased by 2% in women and increased by 5% in men. The median serum 25-hydroxyvitamin D concentration at 12 months was decreased by 19% in women and 10% in men compared with baseline. In the placebo group, this concentration was unchanged in women and increased by 1% in men.

In the study of patients with glucocorticoid-induced osteoporosis, the effects of teriparatide on serum phosphorus were similar to those observed in postmenopausal women with osteoporosis not taking glucocorticoids.

Effects on Markers of Bone Turnover — Daily administration of teriparatide to men and postmenopausal women with osteoporosis in clinical studies stimulated bone formation, as shown by increases in the formation markers serum bone-specific alkaline phosphatase (BSAP) and procollagen 1 carboxy-terminal propeptide (PICP). Data on biochemical markers of bone turnover were available for the first 12 months of treatment. Peak concentrations of PICP at 1 month of treatment were approximately 41% above baseline, followed by a decline to near-baseline values by 12 months. BSAP concentrations increased by 1 month of treatment and continued to rise more slowly from 6 through 12 months. The maximum increases of BSAP were 45% above baseline in women and 23% in men. After discontinuation of therapy, BSAP concentrations returned toward baseline. The increases in formation markers were accompanied by secondary increases in the markers of bone resorption: urinary N-telopeptide (NTX) and urinary deoxyypyridoline (DPP), consistent with the physiological coupling of bone formation and resorption in skeletal remodeling. Changes in BSAP, NTX, and DPP were lower in men than in women, possibly because of lower systemic exposure to teriparatide in men.

In the study of patients with glucocorticoid-induced osteoporosis, the effects of teriparatide on serum markers of bone turnover were similar to those observed in postmenopausal women not taking glucocorticoids.

Digoxin — In a study of 15 healthy people administered digoxin daily to steady state, a single teriparatide dose did not alter the effect of digoxin on the systolic time interval (from electrocardiographic Q-wave onset to aortic valve closure, a measure of digoxin's calcium-mediated cardiac effect). However, sporadic case reports have suggested that hypercalcemia may predispose patients to digitalis toxicity. Because teriparatide may transiently increase serum calcium, teriparatide injection should be used with caution in patients taking digoxin [see *Drug Interactions (7.1)*].

Hydrochlorothiazide — In a study of 20 healthy people, the coadministration of hydrochlorothiazide 25 mg with teriparatide did not affect the serum calcium response to teriparatide 40 mcg. The 24-hour urine excretion of calcium was reduced by a clinically unimportant amount (15%). The effect of coadministration of a higher dose of hydrochlorothiazide with teriparatide on serum calcium levels was not studied [see *Drug Interactions (7.2)*].

Euroside — In a study of 9 healthy people and 17 patients with mild, moderate, or severe renal impairment (CrCl 13 to 72 mL/min), coadministration of intravenous euroside (20 to 100 mcg) with teriparatide 40 mcg resulted in small increases in the serum calcium (2%) and 24-hour urine calcium (37%) responses to teriparatide that did not appear to be clinically important [see *Drug Interactions (7.3)*].

12.3 Pharmacokinetics

The pharmacokinetic parameters of teriparatide following single subcutaneous administration of teriparatide injection in patients with osteoporosis are shown in Table 2.

	Mean ± S.D.	n
T _{max} (hr) ^a	0.25 (0.12, 1.08)	83
C _{max} (pg/mL)	109.5 ± 62.8	83
AUC ₀₋₁ (pg-hr/mL)	134.8 ± 79.7	83
AUC ₀₋₂₄ (pg-hr/mL)	149.8 ± 68.1	72
t _{1/2} (hour)	0.79 ± 0.35	72

Arithmetic mean ± S.D.^a; T_{max} = median (minimum, maximum)

Absorption — Teriparatide is absorbed after subcutaneous injection; the absolute bioavailability is approximately 95% based on plasma data from 20-, 40-, and 80-mcg doses. The rates of absorption and elimination are rapid. The peptide reaches its peak serum concentration about 30 minutes after subcutaneous injection of a 20-mcg dose and declines to non-quantifiable concentrations within 3 hours.

Distribution — Systemic clearance of teriparatide (approximately 62 L/hr in women and 94 L/hr in men) exceeds the rate of normal liver plasma flow, consistent with both hepatic and extra-hepatic clearance. Volume of distribution, following intravenous injection, is approximately 0.12 L/kg. The half-life of teriparatide in serum is 5 minutes when administered by intravenous injection. The longer half-life following subcutaneous administration reflects the time required for absorption from the injection site.

Elimination

Metabolism and Excretion — No metabolism or excretion studies have been performed with teriparatide. However, the mechanisms of metabolism and elimination of PTH(1-34) and intact PTH have been extensively described in published literature. Peripheral metabolism of PTH is believed to occur by non-specific enzymatic mechanisms in the liver followed by excretion via the kidneys.

Specific Populations

Pediatric Patients — Pharmacokinetic data in pediatric patients are not available [see *Warnings and Precautions (5.1)*].

Geriatric Patients — No age-related differences in teriparatide pharmacokinetics were detected (range 31 to 85 years).

Gender — Although systemic exposure to teriparatide was approximately 20% to 30% lower in men than women, the recommended dose for both genders is 20 mcg/day.

Race — The influence of race has not been determined.

Renal Impairment — No pharmacokinetic differences were identified in 11 patients with mild or moderate renal impairment (creatinine clearance (CrCl) 30 to 72 mL/min) administered a single dose of teriparatide. In 5 patients with severe renal impairment (CrCl <30 mL/min), the AUC and T_{1/2} of teriparatide were increased by 73% and 77%, respectively. Maximum serum concentration of teriparatide was not increased. No studies have been performed in patients undergoing dialysis for chronic renal failure [see *Use in Specific Populations (8.7)*].

Hepatic Impairment — No studies have been performed in patients with hepatic impairment [see *Use in Specific Populations (8.6)*].

Drug Interaction Studies

No pharmacokinetic drug-drug interaction studies have been conducted with teriparatide.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis — Two carcinogenicity bioassays were conducted in Fischer 344 rats. In the first study, male and female rats were given daily subcutaneous teriparatide injections of 5, 30, or 75 mcg/kg/day for 24 months from 2 months of age. These doses resulted in systemic exposures that were, respectively, 3, 20, and 60 times higher than the systemic exposure observed in humans following a subcutaneous dose of 20 mcg (based on AUC comparison). Teriparatide treatment resulted in a marked dose-related increase in the incidence of osteosarcoma, a rare malignant bone tumor, in both male and female rats. Osteosarcomas were observed at all doses and the incidence reached 40% to 50% in the high-dose groups. Teriparatide also caused a dose-related increase in osteoblastoma and osteoma in both sexes. No osteosarcomas, osteoblastomas or osteomas were observed in untreated control rats. The bone tumors in rats occurred in association with a large increase in bone mass and focal osteoblast hyperplasia.

The second 2-year study was carried out in order to determine the effect of treatment duration and animal age on the development of bone tumors. Female rats were treated for different periods between 2 and 26 months of age with subcutaneous doses of 5 and 30 mcg/kg (equivalent to 3 and 20 times the human exposure at the 20-mcg dose, based on AUC comparison). The study showed that the occurrence of osteosarcoma, osteoblastoma and osteoma was dependent upon dose and duration of exposure. Bone tumors were observed when immature 2-month old rats were treated with 30 mcg/kg/day for 24 months or with 5 or 30 mcg/kg/day for 6 months. Bone tumors were also observed when mature 6-month old rats were treated with 30 mcg/kg/day for 6 or 20 months. Tumors were not detected when mature 6-month old rats were treated with 5 mcg/kg/day for 6 or 20 months. The results did not demonstrate a difference in susceptibility to bone tumor formation, associated with teriparatide treatment, between mature and immature rats. The relevance of these animal findings to humans is uncertain.

Mutagenesis — Teriparatide was not genotoxic in any of the following test systems: the Ames test for bacterial mutagenesis; the mouse lymphoma assay for mammalian cell mutation; the chromosomal aberration assay in Chinese hamster ovary cells, with and without metabolic activation; and *in vivo* micronucleus test in mice.

Impairment of Fertility — No effects on fertility were observed in male and female rats given subcutaneous teriparatide doses of 30, 100, or 300 mcg/kg/day prior to mating and in females continuing through gestation Day 6 (16 to 160 times the human dose of 20 mcg based on surface area, mcg/m²).

13.2 Animal Toxicology and/or Pharmacology

In single-dose rodent studies using subcutaneous injection of teriparatide, no mortality was seen in rats given doses of 1000 mcg/kg (540 times the human dose based on surface area, mcg/m²) or in mice given 10,000 mcg/kg (2700 times the human dose based on surface area, mcg/m²).

In a long-term study, skeletally mature ovariectomized female monkeys (N=30 per treatment group) were given either daily subcutaneous teriparatide injections of 5 mcg/kg or vehicle. Following the 18-month treatment period, the monkeys were removed from teriparatide treatment and were observed for an additional 3 years. The 5 mcg/kg dose resulted in systemic exposures that were approximately 6 times higher than the systemic exposure observed in humans following a subcutaneous dose of 20 mcg (based on AUC comparison). Bone tumors were not detected by radiographic or histologic evaluation in any monkey in the study.

14 CLINICAL STUDIES

14.1 Treatment of Osteoporosis in Postmenopausal Women

The safety and efficacy of once-daily teriparatide, median exposure of 19 months, were examined in a double-blind, multicenter, placebo-controlled clinical study of 1637 postmenopausal women with osteoporosis (teriparatide 20 mcg, n=541).

All women received 1000 mcg of calcium and at least 400 IU of vitamin D per day. Baseline and endpoint spinal radiographs were evaluated using the semiquantitative scoring. Ninety percent of the women in the study had 1 or more radiographically diagnosed vertebral fractures at baseline. The primary efficacy endpoint was the occurrence of new radiographically diagnosed vertebral fractures defined as changes in the height of previously undeformed vertebrae. Such fractures are not necessarily symptomatic.

Effect on Fracture Incidence

New Vertebral Fractures — Teriparatide, when taken with calcium and vitamin D and compared with calcium and vitamin D alone, reduced the risk of 1 or more new vertebral fractures from 14.3% of women in the placebo group to 5.0% in the teriparatide group. This difference was statistically significant (p <0.001); the absolute reduction in risk was 9.3% and the relative reduction was 65%. Teriparatide was effective in reducing the risk for vertebral fractures regardless of age, baseline rate of bone turnover, or baseline BMD [see *Table 3*].

Table 3. Effect of Teriparatide on Risk of Vertebral Fractures in Postmenopausal Women with Osteoporosis

Percent of Women With Fracture	Teriparatide		Absolute Risk Reduction (%; 95% CI)	Relative Risk Reduction (%; 95% CI)
	(N=444)	(N=448)		
New fracture (≥1)	5.0 ^a	14.3	9.3 (5.5-13.1)	65 (45-78)
1 fracture	3.8	9.4		
2 fractures	0.9	2.9		
≥3 fractures	0.2	2.0		

^a p <0.001 compared with placebo.

New Nonvertebral Osteoporotic Fractures — Teriparatide significantly reduced the risk of any nonvertebral fracture from 5.5% in the placebo group to 2.6% in the teriparatide group (p <0.05). The absolute reduction in risk was 2.9% and the relative reduction was 53%. The incidence of new nonvertebral fractures in the teriparatide group compared with the placebo group was ankle/foot (0.2%, 0.7%), hip (0.2%, 0.7%), humerus (0.4%, 0.4%), pelvis (0.0%, 0.6%), ribs (0.6%, 0.9%), wrist (0.4%, 1.3%), and other sites (1.1%, 1.5%), respectively.

The cumulative percentage of postmenopausal women with osteoporosis who sustained new nonvertebral fractures was lower in women treated with teriparatide than in women treated with placebo (see *Figure 1*).




Figure 1. Cumulative Percentage of Postmenopausal Women with Osteoporosis Sustaining New Nonvertebral Osteoporotic Fractures

Effect on Bone Mineral Density (BMD)

Teriparatide increased lumbar spine BMD in postmenopausal women with osteoporosis. Statistically significant increases were seen at 3 months and continued throughout the treatment period. Postmenopausal women with osteoporosis who were treated with teriparatide had statistically significant increases in BMD from baseline to endpoint at the lumbar spine, femoral neck, total hip, and total body (see *Table 4*).

 Table 4. Mean Percent Change in BMD from Baseline to Endpoint^a in Postmenopausal Women with Osteoporosis, Treated with Teriparatide or Placebo for a Median of 19 Months

	Teriparatide N=541	Placebo N=544
Lumbar spine BMD	9.7 ^b	1.1
Femoral neck BMD	2.8 ^c	-0.7
Total hip BMD	2.6 ^c	-1.0
Trochanter BMD	3.5 ^c	-0.2
Intertrochanter BMD	2.6 ^c	-1.3
Ward's triangle BMD	4.2 ^c	-0.8
Total body BMD	0.6 ^c	-0.5
Distal 1/3 radius BMD	-2.1	-1.3
Ultradistal radius BMD	-0.1	-1.6

^a Intent-to-treat analysis, last observation carried forward.

^b p <0.001 compared with placebo.

^c p <0.05 compared with placebo.

Teriparatide treatment increased lumbar spine BMD from baseline in 96% of postmenopausal women treated. Seventy-two percent of patients treated with teriparat